

# Derivation of cardiac output and alveolar ventilation rate based on energy expenditure measurements in healthy males and females

Pierre Brochu,<sup>a\*</sup> Jules Brodeur<sup>b</sup> and Kannan Krishnan<sup>b</sup>

**ABSTRACT:** Physiologically based pharmacokinetic modeling and occupational exposure assessment studies often use minute ventilation rates ( $VE$ ), alveolar ventilation rates ( $VA$ ) and cardiac outputs ( $Q$ ) that are not reflective of the physiological variations encountered during the aggregate daytime activities of individuals from childhood to adulthood. These variations of  $VE$ ,  $VA$  and  $Q$  values were determined for healthy normal-weight individuals aged 5–96 years by using two types of published individual data that were measured in the same subjects ( $n=902$ ), namely indirect calorimetry measurements and the disappearance rates of oral doses of deuterium ( $^2\text{H}$ ) and heavy-oxygen ( $^{18}\text{O}$ ) in urine monitored by gas-isotope-ratio mass spectrometry. Arteriovenous oxygen content differences ( $0.051\text{--}0.082\text{ ml of O}_2\text{ consumed ml}^{-1}$  of blood) and ratios of the physiological dead space to the tidal volume ( $0.232\text{--}0.419$ ) were determined for oxygen consumption rates ( $0.157\text{--}0.806\text{ l min}^{-1}$ ) required by minute energy expenditures ranging from  $0.76$  to  $3.91\text{ kcal min}^{-1}$ . Generally higher values for the 2.5th up to the 99th percentile for  $VE$  ( $0.132\text{--}0.774\text{ l kg}^{-1}\text{ min}^{-1}$ ,  $4.42\text{--}21.69\text{ l m}^{-2}\text{ min}^{-1}$ ),  $VA$  ( $0.093\text{--}0.553\text{ l kg}^{-1}\text{ min}^{-1}$ ,  $3.09\text{--}15.53\text{ l m}^{-2}\text{ min}^{-1}$ ),  $Q$  ( $0.065\text{--}0.330\text{ l kg}^{-1}\text{ min}^{-1}$ ,  $2.17\text{ to }9.46\text{ l m}^{-2}\text{ min}^{-1}$ ) and ventilation-perfusion ratios ( $1.12\text{--}2.16$ ) were found in children and teenagers aged  $5\text{--}<16.5$  years compared with older individuals. The distributions of cardiopulmonary parameters developed in this study should be useful in facilitating a scientifically sound characterization of the inter-individual differences in the uptake and health risks of lipophilic air pollutants, particularly as they relate to younger children. Copyright © 2011 John Wiley & Sons, Ltd.

**Keywords:** minute energy expenditure; oxygen consumption; minute ventilation; alveolar ventilation; physiological dead space; tidal volume; arteriovenous oxygen content difference; cardiac output; blood flow; ventilation-perfusion ratio; health risk assessment

## INTRODUCTION

In previous publications (Brochu *et al.*, 2006a–c, 2011) we have developed a methodology for the determination of physiological daily inhalation rates of free-living individuals integrating both night-time and daytime respiratory parameters, namely oxygen uptake factors ( $H$ ) and ventilatory equivalents ( $VQ$ ). This approach was based on published input measurements of oxygen consumption rate ( $\text{VO}_2$ ), carbon dioxide production ( $\text{VCO}_2$ ) and minute ventilation rate ( $VE$ ) in a large number of human subjects in order to determine not only the central values but also the standard deviations of  $H$  and  $VQ$  values. The latter values were then integrated with basal daily energy expenditures ( $BEE$ ) and total daily energy expenditures ( $TDEE$ ), that are systematically measured using the doubly labeled water method (DLW), into the calculation process of means and distribution percentiles of physiological daily inhalation rates. This method takes into account voluntary and involuntary energy expended in unrestrained free-living subjects during the entire day (i.e. 24 h), on a daily basis during 7–21 days and only requires periodic body fluid samples (usually urine or saliva) for spectrometric measurements of disappearance rates of oral doses of water isotopes (International Dietary Energy Consultancy Group, 1990).

Physiologically based pharmacokinetic (PBPK) simulation studies allow the determination of the internal dose of xenobiotics. In the case of airborne pollutants, PBPK models require, in addition to many other input parameters, cardiac output and alveolar ventilation rate (Krishnan and Andersen,

2001). PBPK modeling and occupational exposure assessment studies would benefit from the use of values of  $VE$ , alveolar ventilation rate ( $VA$ ) and cardiac output ( $Q$ ) that are reflective of the physiological variations encountered during the aggregate daytime activities over an entire 24 h period, as well as the statistical distribution specific to a group of individuals. For example, the  $VE$  value of  $20.83\text{ l min}^{-1}$ , currently used for occupational exposure assessments, is based on the assumption that workers inhale  $10\text{ m}^3$  in an 8 h workday (US Environmental Protection Agency, 1992). Values for  $VA$  ( $3.83\text{--}5.87\text{ l min}^{-1}$ ) and  $Q$  ( $4.04$  to  $6.73\text{ l min}^{-1}$ ) usually used during PBPK simulation

\*Correspondence to: P. Brochu, Ministère du Développement durable, de l'Environnement et des Parcs, Direction du suivi et de l'état de l'environnement, Service des avis et expertises scientifiques, gouvernement du Québec, édifice Marie-Guyart, 7<sup>e</sup> étage, 675, boulevard René-Lévesque Est, Québec, QC, G1R 5V7, Canada.

E-mail: pierre.brochu@mddep.gouv.qc.ca

<sup>a</sup>Ministère du Développement durable, de l'Environnement et des Parcs, Direction du suivi et de l'état de l'environnement, Service des avis et expertises scientifiques, gouvernement du Québec, édifice Marie-Guyart, 7<sup>e</sup> étage, 675, boulevard René-Lévesque Est, Québec, QC, G1R 5V7, Canada

<sup>b</sup>Département de santé environnementale et santé au travail, Faculté de médecine, Université de Montréal, C.P. 6128, succursale Centre-Ville, Montréal, QC, H3C 3J7, Canada

studies are those for subjects at rest (Arms and Travis, 1988; US Environmental Protection Agency, 1988; Travis and Hattemer-Frey, 1991; Krishnan and Andersen, 2001; Price *et al.*, 2003; Haddad *et al.*, 2006; Valcke and Krishnan, 2009). Finally, despite the fact that variations of minute energy expenditures ( $E$ ) and  $VO_2$  values as a function of time and age are essential for the adequate understanding of the human physiology (Durnin and Passmore, 1967; Elia, 1992, 1997), the distributions of  $E$  and  $VO_2$  percentiles have never been determined from childhood to adulthood. Overall, this may represent a serious shortcoming when establishing indoor or outdoor hygienic standards for airborne toxic chemicals.

The present study is therefore intended to determine the distribution percentiles for  $E$ ,  $VO_2$ ,  $VE$ ,  $Q$  and  $VA$  as a function of age for healthy normal-weight individuals aged 5–96 years during their aggregate daytime activities. In this process, we also developed equations in terms of  $H$ ,  $VQ$ ,  $BEE$  and  $TDEE$  values for converting energy expenditure data into those relevant respiratory and cardiovascular parameters.

## METHODOLOGY

### Study Design

Published  $BEE$  and  $TDEE$  values measured in the same healthy normal-weight individuals aged 5–96 years ( $n = 902$ ) by indirect calorimetry and DLW measurements respectively and taken from the database reported in Institute of Medicine (2002) were converted into  $E$ ,  $VO_2$ ,  $VE$ ,  $Q$ , and  $VA$  values corresponding to their aggregate daytime activities (referred to as  $a$ ). This was done using six types of preliminary parameters integrated into various physiological equations. These include daily energy costs for growth (ECG), sleep duration, the oxygen uptake factor during postprandial phase ( $H_p$ ), arteriovenous oxygen content difference (AVODa), ventilatory equivalents ( $VQa$ ) as well as the ratios of physiological dead space to tidal volume ( $VD_{physa}/VTA$  ratios, unitless). Values for  $BEE$ ,  $TDEE$ , ECG, sleep duration,  $H_p$  and  $VQa$  as well as body weights and heights of subjects per age group are reported in Brochu *et al.* (2011), while values for AVODa and  $VD_{physa}/VTA$  ratios were determined in the present paper. For comparison purposes,  $VAA/Qa$  ratios (unitless) were calculated by using the resulting  $VAA$  and  $Qa$  values. Values for  $Ea$ ,  $VO_2a$ ,  $VEa$ ,  $Qa$  and  $VAA$  were also expressed per unit of body weight and body surface area (BSA in  $m^2$ ). BSA values were calculated using the formula developed by Mosteller (1987) based on height (cm) and body weight (Bw in kg) values:

$$BSA = \left[ \frac{\text{height} \times \text{Bw}}{3600} \right]^{0.5} \quad (1)$$

Some AVODa values and  $VD_{physa}/VTA$  ratios were directly obtained from the literature, but most of them were calculated using published sets of  $Qa$ ,  $VO_2a$  and  $VD_{physa}/VTA$  measurements. In Brochu *et al.* (2011), it was estimated that the oxygenation during aggregate daytime activities ( $VO_2a^*$ ) of males and females aged 5–96 years ranged from 0.18–0.81 and 0.16–0.73  $l \text{ min}^{-1}$ , respectively. Such spans for  $VO_2a^*$  values were used for the adequate selection of input data from the literature. Thus, after the classification of data according to age group, solely the published AVODa,  $Qa$ ,  $VD_{physa}$  and  $VTA$  values and the  $VD_{physa}/VTA$  ratios measured in subjects with experimental  $VO_2$  demands that

were within the span of  $VO_2a^*$  values were included in the present study. These subjects were at rest, in either the sitting or standing position, or performing various activities in the upright position such as exercising on a bicycle ergometer, walking or running on a treadmill or, on a few occasions, performing muscular activities. All published values used in this study were measured at sea level in healthy sedentary untrained and trained individuals with no history of respiratory or cardiac problems when breathing an oxygen concentration of 21%. Data for athletes and explorers were excluded from the calculation process of  $E$ ,  $VO_2$ ,  $VE$ ,  $Q$  and  $VA$  values. Note that children and teenagers are hereafter referred to collectively as children.

### Procedures for Energetic Measurements

The theoretical basis of indirect calorimetry is explained in Ferrannini (1988) and Bursztin *et al.* (1989), while the DLW procedure is discussed at length in International Dietary Energy Consultancy Group (1990). Indirect calorimetry is the most accurate method (Turell and Alexander, 1964) for determining  $BEE$  values based on the equation developed by Weir (1949), where gas exchange (i.e.  $VCO_2$  and  $VO_2$  in  $l \text{ min}^{-1}$ ) is monitored and nitrogen excretion from urine is measured (N in g) in subjects at rest. Values for  $VO_2$  and  $VCO_2$  measured by indirect calorimetry have also been used for the determination of  $H_p$  value by Brochu *et al.* (2011). On the other hand, the DLW method measures the disappearance rates of predetermined oral doses of doubly labeled water ( $^2H_2O$  and  $H_2^{18}O$ ) in free-living subjects, deuterium ( $^2H$ ) and heavy oxygen-18 ( $^{18}O$ ) being monitored in saliva, blood or urine samples by gas-isotope-ratio mass spectrometry over a period of 7–21 consecutive days. Portions of ingested oral doses of  $^2H$  and  $^{18}O$  react with  $CO_2$  to form isotopic carbonic acid which is rapidly transformed into isotopic bicarbonate ions ( $^2HCO_3^-$  and  $HC^{18}OO_2^-$ ) with the catalytic action of carbonic anhydrase. These ions leave erythrocytes to be carried out in the plasma up to the alveolar area. The reverse transformation then occurs in red blood cells where all the  $^2H$  from the  $^2HCO_3^-$  returns to isotopic water molecules ( $^2H_2O$ ), while  $^{18}O$  is returned to the  $H_2^{18}O$ ; some also participate in the formation of isotopic carbon dioxide molecules ( $C^{18}O_2$ ). It is therefore a mixture of non-isotopic ( $CO_2$ ) and isotopic ( $C^{18}O_2$ ) carbon dioxide that is exhaled. The disappearance rate of  $^2H$  reflects water output, while that of  $^{18}O$  represents water output as well as  $VCO_2$  rates. Differences between the two disappearance rates can therefore be used to calculate the  $VCO_2$  rate which is converted into  $TDEE$  values (International Dietary Energy Consultancy Group, 1990).

### Accuracy of Energetic Measurements

Indirect calorimetry measurements of energy expenditure values are accurate within 0.6–0.7% by comparison with those measured by steady-state direct calorimetry in a sealed chamber (or calorimeter) when urinary nitrogen excretions are considered in order to take into account the metabolism of proteins (Turell and Alexander, 1964). However, as do most investigators, the present study avoids the cumbersome correction for the protein metabolism and accepts an error on  $BEE$  values varying from +1 to +2% (Turell and Alexander, 1964) and consequently an error ranging from –2 to –1% on the  $H_p$  value (Brochu *et al.*, 2011). As explained by Brochu *et al.* (2011), the mean precision of  $TDEE$  and ECG values varies from –1.0

to +3.3%. Therefore, the combined effects of, on the one hand, simultaneous mean errors associated with  $H_p$  (i.e. -2 to -1%), BEE (i.e. +1 to +2%), TDEE and ECG (i.e. -1.0 to +3.3%) values on, on the other hand, values of  $VO_2a$ ,  $Qa$ ,  $VEa$ ,  $Va$  were determined in the present study.

### Ea, $VO_2a$ and $VEa$ Values

Precise values for  $VO_2a$  compared with  $VO_2a^*$  ( $l\ min^{-1}$ ) were calculated in this study as well as minute energy expenditures ( $Ea$  in  $kcal\ min^{-1}$ ) and  $VEa$  values ( $l\ min^{-1}$ ). According to Brochu *et al.* (2011), these values can be expressed in terms of BEE, TDEE, ECG ( $kcal$  per day) and sleep duration (Sld in h per day) values by using the following equations:

$$Ea = \left[ \frac{TDEE - BEE}{(24 - Sld) \times 60} \right] + \left[ \frac{BEE + ECG}{1440} \right] \quad (2)$$

$$VO_2a = \left[ \frac{(TDEE - BEE)}{(24 - Sld) \times 60} + \frac{(BEE + ECG)}{1440} \right] \times H_p \quad (3)$$

$$VEa = \left[ \frac{(TDEE - BEE)}{(24 - Sld) \times 60} + \frac{(BEE + ECG)}{1440} \right] \times H_p \times VQa \quad (4)$$

where 1440 and 60 are the conversion factors from days to minutes and hours to minutes, respectively, and 24 is the number of hours in a day.

The value for ECG must be added to BEE in order to take into account the energy requirements for the growth process from birth to adulthood (Brochu *et al.*, 2006a).  $H_p$  is the volume of oxygen consumed (at standard temperature and pressure, dry air, STPD) to produce 1  $kcal$  of energy expended during the postprandial phase.  $VQa$  is the ratio of the  $VEa$  value (at body temperature and saturated with water vapour, BTPS) to the  $VO_2a$  value (at standard temperature and pressure, dry air, STPD), or  $VEa/VO_2a$  ratio (unitless). The value for  $H_p$  of  $0.2059 \pm 0.0019\ l$  of  $O_2\ kcal^{-1}$  ( $n = 1245$ ) and  $VQa$  values varying from  $29.9 \pm 4.2$  to  $32.9 \pm 6.4$  ( $n = 826$ ) according to age group were obtained from Brochu *et al.* (2011).

### Q Values

The Fick principle (Fick, 1870) is one of the cornerstones of human cardiovascular physiology. The physiological mass balance between whole body  $VO_2a$  ( $l\ min^{-1}$ ),  $Qa$  ( $l\ min^{-1}$ ) and the arterial ( $CaO_2$ ) and mixed venous ( $CvO_2$ ) blood oxygen contents ( $ml$  of  $O_2\ ml^{-1}$  of blood), is outlined by the eponymous Fick principle as follows:

$$VO_2a = Qa \times (CaO_2 - CvO_2) = Qa \times AVODa \quad (5)$$

where  $AVODa = O_2$  extraction (i.e. arteriovenous oxygen content difference). Therefore,

$$Qa = \left[ \frac{(TDEE - BEE)}{(24 - Sld) \times 60} + \frac{(BEE + ECG)}{1440} \right] \times \frac{H_p}{AVODa} \quad (6)$$

### VA Values

Values for  $VD_{phys}$  (Bohr, 1891; Enghoff, 1938) include volumes of the conducting airway referred to as anatomical dead space (Fowler, 1948; Folkow and Pappenheimer, 1955) and some

underperfused alveoli (known as the alveolar dead space) not contributing to gas exchange (Guyton, 1991). The  $VA$  is defined as the fraction of the inspired tidal volume per minute ( $VT$  multiplied by the respiratory frequency, known as the  $f$  value) which participates in gas exchange (Guyton, 1991). The  $Va$  ( $l\ min^{-1}$ ) is related to the  $VTa$  ( $l$ ),  $VD_{physa}$  ( $l$ ),  $f$  (number of breaths per minute) and  $VEa$  ( $l\ min^{-1}$ ) values by the following equations (Guyton, 1991):

$$Va = (VTa - VD_{physa}) \times fa \quad (7)$$

$$Va = VEa \times \left[ 1 - \frac{VD_{physa}}{VTa} \right] \quad (8)$$

Therefore,  $Va$  in this study was computed as follows:

$$Va = \left[ \frac{(TDEE - BEE)}{(24 - Sld) \times 60} + \frac{(BEE + ECG)}{1440} \right] \times \left[ 1 - \frac{VD_{physa}}{VTa} \right] \times H_p \times VQa \quad (9)$$

### Sleep Duration

Values for sleep duration in individuals aged 5–96 years ( $n = 13\ 371$ ) taken from Brochu *et al.* (2011) were used in the present study regardless of the proportions of under-, normal-weight, overweight and obese individuals in the cohorts. As showed in Brochu *et al.* (2011), several publications have reported a correlation between sleep curtailment and a higher body mass index (BMI) in children and adults, while others are challenging the view that sleep duration in subjects is inversely related to BMI increases. Therefore, the influence of shorter sleep duration of overweight and obese subjects on the order of magnitude of  $VO_2a$ ,  $Qa$ ,  $VEa$ ,  $Va$  values and  $Va/Qa$  ratios was determined using the calculation process developed by Brochu *et al.* (2011). A first set of data was calculated by using sleep duration reported for a cohort of children aged 7.5–16.5 years (Eisenmann *et al.*, 2006;  $n = 3410$ ) and another of adults 35–74.5 years (Bernsteins *et al.*, 2001;  $n = 6324$ ) for which the proportions of normal-weight, overweight and obese individuals were known. These data were then compared with a second set of values that was calculated when initial sleep durations for 60% of overweight/obese children, and 35% of overweight as well as 55% of obese adults were decreased by 25%. This calculation corresponds to the worst case scenario of sleep duration decrease associated with overweight and obese individuals according to current literature. Further information regarding such a calculation scenario is presented in Brochu *et al.* (2011).

### Statistical Analysis

The best fit distributions (i.e. log-normal or normal) for TDEE, BEE, ECG, sleep duration, body weight, BSA,  $H_p$  and  $VQa$  values have been presented in Brochu *et al.* (2011). Anderson–Darling goodness-of-fit tests were carried out on individual  $Qa$ ,  $AVODa$  and  $Va$  values, as well as  $VD_{physa}/VTa$  ratios from the literature in order to determine their best fit distribution (Cook *et al.*, 1955; Stahlman and Meece, 1957; Johnson *et al.*, 1960; Reeves *et al.*, 1961; Becklake *et al.*, 1962; Donevan *et al.*, 1962; Nelson *et al.*, 1962; Åstrand *et al.*, 1964; Frick and

Somer, 1964; Tabakin *et al.*, 1964; Beaudry *et al.*, 1966; Damato *et al.*, 1966; Ekblom *et al.*, 1968; Ouellet *et al.*, 1969; Hermansen *et al.*, 1970; Jones *et al.*, 1970; Pernow and Saltin, 1971; Frostell *et al.*, 1983; Torre-Bueno *et al.*, 1985).

Means, standard deviations (SD) and distribution percentiles were calculated for AVODa, Ea, VO<sub>2</sub>a, Qa, VEa and VAa values as well as VD<sub>physa</sub>/VTa and VA/Qa ratios. Monte Carlo simulations were conducted based on random sampling involving 10 000 iterations for each calculation process. Distributions were truncated at the minimal and maximal observed values based on a critical analysis of the data compiled from an exhaustive review of the literature. This was done to eliminate from Monte Carlo simulations any outliers that did not remain within the bounds of physiological constraints.

## RESULTS

Mean and SD values as well as distribution percentiles of AVODa, VD<sub>physa</sub>/VTa, Ea, VO<sub>2</sub>a, VEa, Qa, VAa and VAa/Qa for subjects aged 5–96 years are reported in Tables 1–8 respectively. Mean values as a function of age for Ea and VO<sub>2</sub>a, as well as those for VEa, Qa and VAa are presented in Figures 1–3 and 4–6, respectively. Compared with rates expressed per unit of body surface area, those expressed per unit of body weight gradually decrease with age: values for Qa are reduced by 45 and 50%, respectively, while Ea, VO<sub>2</sub>a, VEa, VAa decrease by 51–59% from 5–96 years (Tables 3–7).

Individual Qa (*n* = 129) and AVODa (*n* = 129) values were found to have a better fit with log-normal distributions according to Anderson–Darling goodness-of-fit tests, compared with

**Table 1.** Distribution percentiles of arteriovenous oxygen content differences for aggregate daytime activities of healthy individuals aged 5–96 years

Age group for both gender (years)		Arteriovenous oxygen content differences <sup>a</sup> (ml of O <sub>2</sub> consumed/ml of blood)										
		Percentiles <sup>b</sup>										
	<i>n</i>	Mean ± SD	Min	Max	2.5nd	10th	25th	50th	75th	90th	97.5th	99th
5 to <16.5	110	0.073 ± 0.004	0.057	0.088	0.065	0.067	0.070	0.073	0.075	0.078	0.081	0.082
16.5 to <25	286	0.060 ± 0.005	0.049	0.076	0.051	0.054	0.056	0.060	0.063	0.066	0.070	0.072
25 to <45	193	0.062 ± 0.004	0.048	0.078	0.054	0.057	0.059	0.062	0.064	0.067	0.070	0.072
45 to ≤96	30	0.059 ± 0.003	0.051	0.069	0.054	0.056	0.057	0.059	0.061	0.063	0.065	0.066

<sup>a</sup>Measurements reported in Johnson *et al.* (1960), Reeves *et al.* (1961), Donevan *et al.* (1962), Åstrand *et al.* (1964), Frick and Somer (1964), Tabakin *et al.* (1964), Dagenais *et al.* (1966), Damato *et al.* (1966), Ekblom *et al.* (1968), Ouellet *et al.* (1969), Hermansen *et al.* (1970), Jones *et al.* (1970), Eriksson *et al.* (1971), Pernow and Saltin (1971), Krone *et al.* (1972), Zeidifard *et al.* (1972), Sharma *et al.* (1977), Kanstrup and Ekblom (1978), Hossack and Bruce (1982), Frostell *et al.* (1983), Lewis *et al.* (1983), Torre-Bueno *et al.* (1985), Wagner *et al.* (1986), Bebout *et al.* (1989), Miyamoto *et al.* (1989), Podolsky *et al.* (1996), Turley and Wilmore (1997a), Rice *et al.* (1999), Hopkins *et al.* (2000), Sun *et al.* (2000), McGuire *et al.* (2001), Nottin *et al.* (2002), Poole *et al.* (2002), Vinet *et al.* (2002), Gisolf *et al.* (2003), Olfert *et al.* (2004), Dibski *et al.* (2005). <sup>b</sup>Percentiles based on a log-normal distribution according to the Anderson–Darling test performed on individual data. *n* = number of individuals; SD = standard deviation; Min = minimum; Max = maximum.

**Table 2.** Distribution percentiles of ratios of physiological dead space to tidal volume for aggregate daytime activities of healthy individuals aged 5–96 years

Age group for both gender (years)		Ratios of physiological dead space to tidal volume <sup>a</sup> (unitless)										
		Percentiles <sup>b</sup>										
		<i>n</i>	Mean ± SD	Min	Max	2.5nd	10th	25th	50th	75th	90th	97.5th
5 to <10 <sup>c</sup>	52	0.336 ± 0.040	0.244	0.428	0.264	0.287	0.311	0.337	0.363	0.398	0.409	0.418
10 to <16.5 <sup>d</sup>	81	0.294 ± 0.032	0.203	0.386	0.232	0.253	0.272	0.293	0.315	0.345	0.357	0.366
16.5 to <25 <sup>e</sup>	48	0.301 ± 0.026	0.220	0.386	0.250	0.268	0.283	0.300	0.318	0.343	0.351	0.361
25 to <35 <sup>f</sup>	112	0.329 ± 0.015	0.280	0.389	0.299	0.310	0.319	0.329	0.339	0.354	0.359	0.364
35 to <45 <sup>g</sup>	79	0.344 ± 0.018	0.281	0.407	0.308	0.321	0.331	0.344	0.356	0.374	0.380	0.386
45 to <65 <sup>h</sup>	55	0.339 ± 0.021	0.273	0.405	0.299	0.312	0.325	0.340	0.354	0.374	0.380	0.388
65 to ≤96 <sup>i</sup>	36	0.381 ± 0.018	0.334	0.428	0.347	0.359	0.369	0.382	0.393	0.410	0.414	0.419

<sup>a</sup>VDphysa/VTa ratios. <sup>b</sup>Percentiles based on a normal distribution according to the Anderson–Darling test performed on individual data. <sup>c</sup>Kerr (1976). <sup>d</sup>Beaudry *et al.* (1966) and Kerr (1976).<sup>e</sup>Mellemgaard (1966), Whipp and Wasserman (1969), Olfert *et al.* (2004). <sup>f</sup>Froeb (1962), Malmberg (1966), Mellemgaard (1966), Whipp and Wasserman (1969), Craig *et al.* (1971), Frostell *et al.* (1983), Allen *et al.* (1984), Dempsey *et al.* (1984), Olfert *et al.* (2004). <sup>g</sup>Froeb (1962), Malmberg (1966), Mellemgaard (1966), Craig *et al.* (1971), Dempsey *et al.* (1984). <sup>h</sup>Mellemgaard (1966), Craig *et al.* (1971), Frostell *et al.* (1983). <sup>i</sup>Tenney and Miller (1956), Mellemgaard (1966), Craig *et al.* (1971). *n* = number of individuals; SD = standard deviation; Min = minimum; Max = maximum.



Minute energy expenditures<sup>a</sup>

Age group (years)	Males								Females								
	Mean ± SD				Percentiles				Mean ± SD				Percentiles				
	2.5nd	10th	25th	50th	75th	90th	97.5th	99th	2.5nd	10th	25th	50th	75th	90th	97.5th	99th	
								(kcal min <sup>-1</sup> )									
5 to <7	1.22±0.24	0.81	0.92	1.05	1.21	1.37	1.54	1.75	1.85	1.18±0.22	0.79	0.89	1.01	1.16	1.32	1.48	1.66
7 to <10	1.54±0.36	0.96	1.10	1.27	1.50	1.76	2.03	2.36	2.54	1.48±0.31	0.97	1.11	1.26	1.46	1.68	1.90	2.17
10 to <16.5	2.27±0.63	1.29	1.50	1.78	2.18	2.68	3.16	3.70	3.91	1.93±0.50	1.12	1.32	1.56	1.87	2.25	2.61	3.05
16.5 to <25	2.68±0.45	1.83	2.07	2.35	2.69	3.02	3.27	3.49	3.64	2.22±0.32	1.62	1.81	2.00	2.21	2.44	2.63	2.86
25 to <35	2.55±0.48	1.77	1.97	2.20	2.50	2.85	3.20	3.63	3.86	2.06±0.41	1.36	1.56	1.76	2.03	2.32	2.60	2.96
35 to <45	2.49±0.33	1.92	2.07	2.24	2.48	2.72	2.94	3.14	3.23	2.08±0.29	1.56	1.69	1.85	2.07	2.29	2.47	2.64
45 to <65	2.25±0.42	1.57	1.72	1.93	2.20	2.54	2.84	3.15	3.27	1.82±0.34	1.17	1.37	1.57	1.81	2.05	2.28	2.52
65 to ≤96	1.89±0.44	1.14	1.34	1.57	1.86	2.19	2.50	2.84	2.97	1.41±0.39	0.76	0.92	1.12	1.37	1.65	1.96	2.27
									(kcal kg <sup>-1</sup> min <sup>-1</sup> ) <sup>b</sup>								
5 to <7	0.063±0.014	0.040	0.046	0.053	0.061	0.071	0.081	0.093	0.101	0.061±0.013	0.038	0.044	0.051	0.059	0.069	0.079	0.090
7 to <10	0.059±0.017	0.033	0.040	0.047	0.057	0.068	0.082	0.098	0.108	0.055±0.013	0.033	0.039	0.045	0.053	0.063	0.073	0.084
10 to <16.5	0.055±0.020	0.026	0.033	0.040	0.052	0.067	0.083	0.103	0.116	0.044±0.014	0.023	0.028	0.034	0.042	0.053	0.063	0.078
16.5 to <25	0.038±0.007	0.025	0.029	0.033	0.038	0.043	0.047	0.052	0.055	0.037±0.006	0.026	0.029	0.033	0.037	0.041	0.045	0.050
25 to <35	0.036±0.008	0.024	0.027	0.031	0.035	0.040	0.046	0.053	0.057	0.036±0.008	0.022	0.026	0.030	0.035	0.041	0.046	0.053
35 to <45	0.036±0.006	0.026	0.029	0.032	0.035	0.039	0.043	0.048	0.050	0.035±0.006	0.026	0.028	0.031	0.035	0.039	0.043	0.047
45 to <65	0.032±0.007	0.021	0.024	0.027	0.031	0.036	0.041	0.046	0.049	0.031±0.006	0.020	0.023	0.026	0.031	0.035	0.039	0.044
65 to ≤96	0.028±0.007	0.016	0.019	0.023	0.027	0.032	0.037	0.043	0.046	0.025±0.007	0.013	0.016	0.019	0.024	0.029	0.035	0.042
									(kcal m <sup>-2</sup> min <sup>-1</sup> ) <sup>b</sup>								

<sup>a</sup> $E_a = [(TDEE - BEE)/(24 - Sld) \times 60] + (BEE + ECG/1440)$ , where TDEE = total daily energy expenditure, BEE = basal energy expenditure and ECG = stored daily energy cost for growth. <sup>b</sup> $E_a$  (kcal min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in kcal kg<sup>-1</sup> min<sup>-1</sup> and kcal m<sup>-2</sup> min<sup>-1</sup>, respectively. <sup>a</sup>Values for TDEE, BEE, ECG (kcal per day), Sld (h per day), Bw (kg) and BSA (m<sup>2</sup>) are presented in Brochu *et al.* (2011). SD = standard deviation.

Oxygen consumption rates<sup>a</sup>[illegible]

<sup>a</sup> $\text{VO}_2\alpha = [(TDEE - BEE) / (24 - \text{Sld}) \times 60] + (BEE + \text{ECG}) / 1440 \times H_p$ , where  $H_p$  = oxygen uptake factor. TDEE, BEE, ECG and Sld are defined in Table 3. <sup>b</sup> $\text{VO}_2\alpha$  ( $\text{L min}^{-1}$ ) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in  $\text{l kg}^{-1} \text{min}^{-1}$  and  $\text{l m}^{-2} \text{min}^{-1}$ , respectively. <sup>a</sup>, <sup>b</sup>Values for TDEE, BEE, ECG (kcal per day), Sld (h per day), Bw (kg), BSA ( $\text{m}^2$ ) and HP ( $0.2059 \pm 0.0019$  l of  $\text{O}_2 \text{kcal}^{-1}$ ) are reported in Brochu *et al.* (2011). SD = standard deviation.

**Table 5.** Distribution percentiles of minute ventilation rates for aggregate daytime activities of normal-weight individuals aged 5–96 years

Age group (years)	Minute ventilation rates <sup>a</sup>										Females				
	Males					Mean ± SD					Percentiles				
	2.5nd	10th	25th	50th	75th	90th	97.5th	99th	(l min <sup>-1</sup> )		2.5th	10th	25th	50th	75th
5 to <7	7.75 ± 1.55	5.10	5.82	6.63	7.64	8.75	11.10	11.84	7.45 ± 1.44		6.41	5.65	6.41	7.32	8.36
7 to <10	9.74 ± 2.32	6.01	6.94	8.02	9.50	11.15	15.01	16.17	9.40 ± 1.96		7.98	7.00	7.98	9.22	10.64
10 to <16.5	13.94 ± 4.42	7.28	8.82	10.68	13.25	16.52	24.20	26.51	11.87 ± 3.55		9.27	7.70	9.27	11.40	13.98
16.5 to <25	17.91 ± 4.54	10.63	12.51	14.61	17.35	20.63	28.04	30.41	14.83 ± 3.48		12.33	10.73	12.33	14.40	16.91
25 to <35	17.17 ± 4.12	10.69	12.34	14.19	16.64	19.64	26.57	29.04	13.86 ± 3.43		11.42	9.82	11.42	13.51	15.84
35 to <45	16.95 ± 4.94	9.42	11.25	13.39	16.20	19.83	23.55	28.27	14.15 ± 4.26		11.07	9.28	11.07	13.52	16.60
45 to <65	15.47 ± 4.40	8.64	10.41	12.29	14.86	17.95	21.39	25.77	12.51 ± 3.59		9.91	8.30	9.91	12.08	14.62
65 to ≤96	13.05 ± 4.17	6.55	8.23	10.03	12.45	15.41	18.63	22.95	9.69 ± 3.43		7.16	5.72	7.16	9.14	11.65
									(l kg <sup>-1</sup> min <sup>-1</sup> ) <sup>b</sup>						
5 to <7	0.397 ± 0.089	0.249	0.289	0.333	0.389	0.450	0.515	0.592	0.645	0.383 ± 0.086	0.241	0.280	0.322	0.375	0.436
7 to <10	0.374 ± 0.107	0.209	0.251	0.296	0.359	0.435	0.520	0.623	0.693	0.348 ± 0.084	0.209	0.246	0.286	0.339	0.400
10 to <16.5	0.341 ± 0.135	0.151	0.193	0.242	0.316	0.413	0.517	0.675	0.774	0.273 ± 0.096	0.132	0.166	0.203	0.259	0.324
16.5 to <25	0.255 ± 0.068	0.148	0.175	0.205	0.246	0.295	0.347	0.412	0.442	0.248 ± 0.062	0.149	0.175	0.203	0.240	0.284
25 to <35	0.242 ± 0.062	0.145	0.170	0.197	0.233	0.279	0.325	0.386	0.422	0.240 ± 0.065	0.137	0.163	0.193	0.231	0.277
35 to <45	0.244 ± 0.074	0.132	0.159	0.191	0.232	0.286	0.343	0.415	0.455	0.242 ± 0.076	0.128	0.157	0.187	0.230	0.284
45 to <65	0.217 ± 0.065	0.118	0.143	0.170	0.208	0.254	0.303	0.372	0.414	0.214 ± 0.063	0.114	0.140	0.168	0.206	0.251
65 to ≤96	0.193 ± 0.064	0.096	0.119	0.146	0.184	0.227	0.278	0.349	0.385	0.171 ± 0.064	0.076	0.098	0.124	0.161	0.206
									(l m <sup>-2</sup> min <sup>-1</sup> ) <sup>b</sup>						
5 to <7	9.88 ± 2.10	6.35	7.34	8.37	9.68	11.21	12.61	14.54	15.57	9.53 ± 1.98	6.18	7.08	8.11	9.34	10.75
7 to <10	10.10 ± 2.62	5.91	6.99	8.19	9.78	11.65	13.64	16.12	17.54	9.51 ± 2.12	5.97	6.97	7.98	9.32	10.85
10 to <16.5	10.56 ± 3.77	5.04	6.28	7.78	9.95	12.67	15.65	19.53	21.69	8.61 ± 2.76	4.42	5.43	6.57	8.19	10.21
16.5 to <25	9.60 ± 2.49	5.58	6.68	7.79	9.28	11.09	12.90	15.21	16.54	8.85 ± 2.13	5.42	6.34	7.29	8.62	10.15
25 to <35	9.15 ± 2.27	5.56	6.49	7.51	8.84	10.49	12.14	14.36	15.77	8.52 ± 2.19	4.95	5.96	6.95	8.29	9.78
35 to <45	9.19 ± 2.71	5.07	6.06	7.22	8.78	10.72	12.83	15.55	16.99	8.64 ± 2.63	4.62	5.65	6.73	8.26	10.14
45 to <65	8.30 ± 2.41	4.61	5.50	6.55	7.97	9.66	11.52	13.91	15.46	7.68 ± 2.23	4.12	5.08	6.07	7.40	9.00
65 to ≤96	7.21 ± 2.35	3.60	4.51	5.50	6.87	8.50	10.36	12.82	14.21	6.08 ± 2.20	2.80	3.54	4.47	5.75	7.32

<sup>a</sup>VEa = [(TDEE - BEE)/(24 - Sld) × 60] + (BEE + ECG/1440) × Hp × VQa, where Hp = ventilatory uptake factor and VQa = ventilatory equivalent. TDEE, BEE, ECG and Sld are defined in Table 3. <sup>b</sup>VEa (l min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in l kg<sup>-1</sup> min<sup>-1</sup> and l m<sup>-2</sup> min<sup>-1</sup> respectively. <sup>a</sup>, <sup>b</sup>Values for TDEE, BEE, ECG (kcal per day), Sld (h per day), Bw (kg), BSA (m<sup>2</sup>), Hp (0.2059 ± 0.0019 l of O<sub>2</sub> kcal<sup>-1</sup>) and VQa (unitless) are given in Brochu et al. (2011). SD = standard deviation.

**Table 6.** Distribution percentiles of cardiac outputs for aggregate daytime activities of normal-weight individuals aged 5–96 years

Age group (years)	Cardiac outputs <sup>a</sup>										Males					Females						
	Mean ± SD					Percentiles					Mean ± SD					Percentiles						
	2.5nd	10th	25th	50th	75th	90th	97.5th	99th	(l min <sup>-1</sup> )					2.5nd	10th	25th	50th	75th	90th	97.5th	99th	
5 to <7	3.48 ± 0.71	2.28	2.60	2.96	3.42	3.93	4.43	5.04	5.40	3.35 ± 0.66					2.23	2.53	2.86	3.31	3.77	4.23	4.77	5.05
7 to <10	4.35 ± 1.04	2.69	3.10	3.59	4.24	4.98	5.77	6.73	7.22	4.22 ± 0.91					2.70	3.10	3.55	4.13	4.78	5.45	6.23	6.67
10 to <16.5	6.44 ± 1.83	3.61	4.25	5.04	6.18	7.60	9.03	10.52	11.19	5.48 ± 1.45					3.11	3.72	4.41	5.33	6.39	7.45	8.78	9.48
16.5 to <25	9.27 ± 1.71	6.14	7.02	8.01	9.23	10.46	11.53	12.66	13.23	7.68 ± 1.25					5.45	6.11	6.79	7.61	8.51	9.32	10.28	10.87
25 to <35	8.56 ± 1.71	5.80	6.50	7.29	8.38	9.59	10.86	12.40	13.19	6.91 ± 1.46					4.43	5.13	5.88	6.79	7.79	8.84	10.12	10.89
35 to <45	8.35 ± 1.22	6.21	6.81	7.44	8.28	9.17	10.02	10.89	11.35	6.97 ± 1.08					5.07	5.59	6.17	6.91	7.70	8.40	9.17	9.55
45 to <65	7.85 ± 1.51	5.39	5.96	6.71	7.68	8.84	9.97	11.08	11.65	6.34 ± 1.24					4.02	4.75	5.47	6.31	7.17	7.99	8.89	9.34
65 to ≤96	6.61 ± 1.57	3.95	4.67	5.43	6.48	7.66	8.79	9.96	10.54	4.91 ± 1.39					2.64	3.21	3.89	4.75	5.78	6.84	8.01	8.52
5 to <7	0.178 ± 0.041	0.111	0.129	0.149	0.174	0.203	0.233	0.269	0.290	0.172 ± 0.039					0.107	0.125	0.143	0.168	0.196	0.223	0.258	0.279
7 to <10	0.167 ± 0.048	0.094	0.112	0.133	0.161	0.194	0.231	0.276	0.305	0.155 ± 0.039					0.092	0.109	0.127	0.151	0.179	0.208	0.244	0.263
10 to <16.5	0.157 ± 0.058	0.073	0.092	0.115	0.148	0.190	0.237	0.296	0.330	0.126 ± 0.040					0.065	0.079	0.096	0.120	0.149	0.179	0.223	0.247
16.5 to <25	0.132 ± 0.027	0.085	0.098	0.113	0.131	0.149	0.167	0.188	0.199	0.128 ± 0.024					0.087	0.099	0.112	0.127	0.143	0.159	0.180	0.192
25 to <35	0.121 ± 0.027	0.078	0.090	0.101	0.117	0.136	0.156	0.180	0.194	0.120 ± 0.029					0.073	0.085	0.099	0.117	0.137	0.157	0.183	0.201
35 to <45	0.120 ± 0.020	0.086	0.095	0.105	0.118	0.133	0.147	0.163	0.172	0.119 ± 0.020					0.084	0.094	0.104	0.118	0.132	0.146	0.162	0.170
45 to <65	0.110 ± 0.024	0.072	0.082	0.093	0.108	0.125	0.143	0.162	0.172	0.108 ± 0.023					0.068	0.079	0.092	0.107	0.123	0.139	0.157	0.166
65 to ≤96	0.098 ± 0.025	0.056	0.067	0.079	0.095	0.113	0.131	0.152	0.163	0.086 ± 0.026					0.045	0.055	0.067	0.083	0.102	0.122	0.147	0.159
5 to <7	4.43 ± 0.96	2.82	3.26	3.74	4.35	5.03	5.71	6.57	7.01	4.29 ± 0.91					2.76	3.16	3.61	4.21	4.86	5.51	6.31	6.72
7 to <10	4.52 ± 1.18	2.66	3.14	3.66	4.38	5.23	6.12	7.20	7.90	4.26 ± 0.98					2.66	3.08	3.55	4.17	4.85	5.57	6.50	6.93
10 to <16.5	4.88 ± 1.60	2.49	3.02	3.68	4.63	5.82	7.10	8.57	9.46	3.97 ± 1.15					2.17	2.62	3.13	3.82	4.67	5.52	6.60	7.21
16.5 to <25	4.97 ± 0.95	3.25	3.74	4.27	4.94	5.61	6.21	6.87	7.24	4.59 ± 0.79					3.18	3.60	4.03	4.54	5.11	5.62	6.25	6.58
25 to <35	4.56 ± 0.95	3.02	3.42	3.87	4.45	5.13	5.85	6.72	7.19	4.25 ± 0.94					2.65	3.12	3.58	4.16	4.81	5.49	6.31	6.86
35 to <45	4.53 ± 0.70	3.32	3.64	4.01	4.49	4.99	5.47	5.98	6.31	4.25 ± 0.69					3.05	3.39	3.75	4.21	4.72	5.17	5.66	5.93
45 to <65	4.21 ± 0.85	2.83	3.16	3.57	4.11	4.77	5.40	6.06	6.41	3.89 ± 0.79					2.45	2.89	3.34	3.86	4.42	4.94	5.51	5.80
65 to ≤96	3.65 ± 0.90	2.14	2.55	2.98	3.57	4.23	4.86	5.60	5.97	3.08 ± 0.90					1.63	2.00	2.42	2.98	3.65	4.31	5.10	5.53

<sup>a</sup>Qa = [(TDEE – BEE)/((24 – Sld) × 60)] + (BEE + ECG)/(1440) × Hp × AVODa<sup>-1</sup>, where Hp = oxygen uptake factor, AVODa = arteriovenous oxygen content differences (ml of O<sub>2</sub> consumed ml<sup>-1</sup> of blood). Values for AVODa are given in Table 1. TDEE, BEE, ECG and Sld are defined in Table 3. <sup>b</sup>Qa (L min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in l kg<sup>-1</sup> min<sup>-1</sup> and l m<sup>-2</sup> min<sup>-1</sup> respectively. <sup>a</sup>, <sup>b</sup>Values for TDEE, BEE, ECG (kcal per day), Bw (kg), BSA (m<sup>2</sup>) and, Hp (0.2059 ± 0.0019 l of O<sub>2</sub> kcal<sup>-1</sup>) were taken from Brochu *et al.* (2011). SD = standard deviation.

<sup>a</sup> $Qa = [(TDEE - BEE)/(24 - Sld) \times 60] + (BEE + ECG/1440) \times H_p \times AVODa^{-1}$ , where  $H_p$  = oxygen uptake factor,  $AVODa$  = arteriovenous oxygen content differences (ml of  $O_2$  consumed ml<sup>-1</sup> of blood). Values for  $AVODa$  are given in Table 1.  $TDEE$ ,  $BEE$ ,  $ECG$  and  $Sld$  are defined in Table 3. <sup>b</sup> $Qa$  (L min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in l kg<sup>-1</sup> min<sup>-1</sup> and l m<sup>-2</sup> min<sup>-1</sup> respectively. <sup>a</sup>, <sup>b</sup>Values for  $TDEE$ ,  $BEE$ ,  $ECG$  (kcal per day),  $Sld$  (h per day),  $Bw$  (kg),  $BSA$  (m<sup>2</sup>) and,  $H_p$  (0.2059 ± 0.0019 l of  $O_2$  kcal<sup>-1</sup>) were taken from Brochu et al. (2011).  $SD$  = standard deviation.



**Table 7.** Distribution percentiles of alveolar ventilation rates for aggregate daytime activities of normal-weight individuals aged 5–96 years

Age group (years)	Alveolar ventilation rates <sup>a</sup>																		
	Males					Females													
	Mean ± SD	Percentiles				Mean ± SD	Percentiles												
		2.5th	10th	25th	50th	75th	90th	97.5th	99th	(l min <sup>-1</sup> )		2.5nd	10th	25th	50th	75th	90th	97.5th	99th
5 to <7	5.14 ± 1.07	3.33	3.83	4.36	5.06	5.82	6.56	7.48	8.02	4.94 ± 1.00	3.25	3.71	4.22	4.86	5.58	6.28	7.13	7.55	
7 to <10	6.47 ± 1.58	3.95	4.58	5.31	6.28	7.41	8.63	10.02	10.93	6.24 ± 1.35	3.99	4.59	5.26	6.11	7.07	8.07	9.28	9.89	
10 to <16.5	9.84 ± 3.15	5.16	6.19	7.51	9.37	11.67	14.11	17.21	18.81	8.38 ± 2.53	4.45	5.41	6.55	8.05	9.85	11.79	14.27	15.77	
16.5 to <25	12.52 ± 3.22	7.38	8.71	10.20	12.11	14.44	16.86	19.76	21.50	10.37 ± 2.47	6.40	7.46	8.58	10.08	11.85	13.64	15.87	17.26	
25 to <35	11.52 ± 2.78	7.15	8.27	9.52	11.17	13.18	15.22	17.82	19.62	9.30 ± 2.30	5.48	6.57	7.67	9.06	10.61	12.38	14.59	15.86	
35 to <45	11.12 ± 3.26	6.16	7.36	8.76	10.64	13.04	15.49	18.69	20.52	9.29 ± 2.81	5.05	6.08	7.25	8.86	10.87	13.06	15.76	17.58	
45 to <65	10.23 ± 2.93	5.68	6.83	8.12	9.83	11.87	14.22	17.12	18.71	8.27 ± 2.39	4.46	5.47	6.53	7.98	9.67	11.48	13.76	15.04	
65 to ≤96	8.07 ± 2.58	4.06	5.08	6.20	7.72	9.54	11.53	14.22	15.50	5.99 ± 2.13	2.80	3.54	4.42	5.67	7.22	8.91	10.96	12.17	
									(l kg <sup>-1</sup> min <sup>-1</sup> ) <sup>b</sup>										
5 to <7	0.263 ± 0.061	0.163	0.191	0.219	0.257	0.300	0.343	0.398	0.435	0.254 ± 0.059	0.157	0.184	0.212	0.248	0.290	0.334	0.386	0.414	
7 to <10	0.248 ± 0.072	0.137	0.165	0.196	0.238	0.289	0.345	0.415	0.464	0.231 ± 0.058	0.138	0.161	0.189	0.224	0.266	0.307	0.361	0.392	
10 to <16.5	0.241 ± 0.096	0.106	0.136	0.170	0.223	0.292	0.366	0.477	0.553	0.193 ± 0.068	0.093	0.116	0.143	0.183	0.229	0.285	0.355	0.395	
16.5 to <25	0.178 ± 0.048	0.103	0.122	0.143	0.172	0.206	0.244	0.290	0.312	0.173 ± 0.044	0.104	0.122	0.141	0.168	0.199	0.233	0.274	0.299	
25 to <35	0.162 ± 0.042	0.097	0.114	0.132	0.156	0.187	0.218	0.259	0.284	0.161 ± 0.044	0.092	0.110	0.129	0.155	0.186	0.219	0.261	0.285	
35 to <45	0.160 ± 0.049	0.086	0.104	0.125	0.152	0.188	0.225	0.273	0.300	0.159 ± 0.050	0.084	0.103	0.123	0.151	0.186	0.226	0.273	0.305	
45 to <65	0.144 ± 0.043	0.077	0.094	0.112	0.138	0.168	0.201	0.246	0.274	0.141 ± 0.042	0.075	0.092	0.111	0.136	0.166	0.198	0.237	0.266	
65 to ≤96	0.119 ± 0.040	0.059	0.074	0.091	0.114	0.141	0.172	0.216	0.237	0.106 ± 0.040	0.047	0.061	0.077	0.100	0.128	0.160	0.200	0.221	
									(l m <sup>-2</sup> min <sup>-1</sup> ) <sup>b</sup>										
5 to <7	6.56 ± 1.44	4.15	4.80	5.52	6.43	7.45	8.47	9.73	10.51	6.32 ± 1.36	4.02	4.65	5.33	6.20	7.16	8.14	9.33	10.09	
7 to <10	6.70 ± 1.78	3.90	4.62	5.40	6.48	7.74	9.08	10.72	11.86	6.31 ± 1.45	3.92	4.57	5.26	6.16	7.20	8.29	9.51	10.31	
10 to <16.5	7.45 ± 2.68	3.54	4.43	5.48	7.03	8.95	11.04	13.82	15.53	6.07 ± 1.96	3.09	3.81	4.63	5.78	7.19	8.68	10.70	11.92	
16.5 to <25	6.71 ± 1.76	3.88	4.66	5.43	6.47	7.75	9.05	10.73	11.63	6.19 ± 1.51	3.77	4.42	5.09	6.02	7.10	8.23	9.54	10.46	
25 to <35	6.14 ± 1.53	3.73	4.35	5.03	5.94	7.05	8.15	9.66	10.57	5.72 ± 1.47	3.30	3.99	4.66	5.56	6.58	7.67	9.03	9.89	
35 to <45	6.03 ± 1.79	3.31	3.97	4.73	5.76	7.06	8.42	10.25	11.23	5.67 ± 1.73	3.02	3.69	4.41	5.41	6.64	7.99	9.65	10.68	
45 to <65	5.49 ± 1.61	3.02	3.63	4.32	5.27	6.39	7.63	9.30	10.20	5.07 ± 1.48	2.71	3.34	4.00	4.89	5.96	7.05	8.51	9.35	
65 to ≤96	4.46 ± 1.45	2.22	2.79	3.42	4.25	5.27	6.40	7.92	8.75	3.76 ± 1.37	1.73	2.19	2.76	3.56	4.54	5.60	6.96	7.74	


<sup>a</sup>V<sub>A</sub>Q = [(TDEE – BEE)/((24 – Sld) × 60) + (BEE + ECG)/(1440)] × H<sub>p</sub> × VQ<sub>a</sub> × (1 – VDphysa/VT<sub>a</sub>), where VDphysa/VT<sub>a</sub> = ratio of the physiological dead space to the tidal volume, H<sub>p</sub> = oxygen uptake factor and VQ<sub>a</sub> = ventilatory equivalent. Values for VDphysa/VT<sub>a</sub> (unitless) are given in Table 2. TDEE, BEE, ECG and Sld are defined in Table 3. <sup>b</sup>V<sub>A</sub>Q (l min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in l kg<sup>-1</sup> min<sup>-1</sup> and l m<sup>-2</sup> min<sup>-1</sup> respectively. <sup>a</sup>, <sup>b</sup>Values for TDEE, BEE, ECG (kcal per day), Sld (h per day), Bw (kg), BSA (m<sup>2</sup>), H<sub>p</sub> (0.2059 ± 0.0019 l of O<sub>2</sub> kcal<sup>-1</sup>) and VQ<sub>a</sub> (unitless) are reported in Brochu *et al.* (2011). SD = standard deviation.

<sup>a</sup> $V_{Aa} = [(TDEE - BEE)/(24 - Sld) \times 60] + (BEE + ECG)/(1440) \times H_p \times VQa \times (1 - VDphysa/VTa)$ , where  $VDphysa/VTa$  = ratio of the physiological dead space to the tidal volume,  $H_p$  = oxygen uptake factor and  $VQa$  = ventilatory equivalent. Values for  $VDphysa/VTa$  (unitless) are given in Table 2. TDEE, BEE, ECG and Sld are defined in Table 3. <sup>b</sup> $V_{Aa}$  (l min<sup>-1</sup>) were divided by body weights (Bw) and body surface areas (BSA) in order to obtain values expressed in l kg<sup>-1</sup> min<sup>-1</sup> and l m<sup>-2</sup> min<sup>-1</sup> respectively. <sup>a</sup>, <sup>b</sup>Values for TDEE, BEE, ECG (kcal per day), Sld (h per day), Bw (kg), BSA (m<sup>2</sup>),  $H_p$  (0.2059 ± 0.0019 l of O<sub>2</sub> kcal<sup>-1</sup>) and  $VQa$  (unitless) are reported in Brochu et al. (2011). SD = standard deviation.

**Table 8.** Distribution percentiles of ventilation-perfusion ratios for aggregate daytime activities of normal-weight individuals aged 5 to 96 years

Age group for both genders (years)	Mean $\pm$ SD	Ventilation-perfusion ratios <sup>a</sup> (l of alveolar air per l of blood)							
		Percentiles							
		2.5nd	10th	25th	50th	75th	90th	97.5th	99th
5 to <7	1.49 $\pm$ 0.12	1.26	1.34	1.40	1.49	1.57	1.64	1.73	1.78
7 to <10	1.49 $\pm$ 0.12	1.26	1.34	1.40	1.49	1.57	1.64	1.73	1.78
10 to <16.5	1.53 $\pm$ 0.24	1.12	1.24	1.37	1.51	1.68	1.84	2.05	2.16
16.5 to <25	1.36 $\pm$ 0.28	0.91	1.03	1.16	1.33	1.52	1.72	1.98	2.14
25 to <35	1.35 $\pm$ 0.22	0.98	1.09	1.20	1.34	1.49	1.63	1.82	1.93
35 to <45	1.34 $\pm$ 0.36	0.76	0.92	1.08	1.29	1.54	1.82	2.16	2.40
45 to <65	1.31 $\pm$ 0.29	0.83	0.97	1.10	1.27	1.48	1.70	1.96	2.11
65 to $\leq$ 96	1.22 $\pm$ 0.27	0.78	0.91	1.03	1.19	1.38	1.58	1.83	1.98

<sup>a</sup>VAa/Qa. Values for Qa (l of blood min<sup>-1</sup>) and VAa (l of alveolar air min<sup>-1</sup>) are given in Tables 6 and 7 respectively. SD = standard deviation.



Plotted values are for midpoint ages of the age cohorts reported in Tables 3 and 4.  
 E = minute energy expenditure rate; VO<sub>2</sub> = oxygen consumption rate; males = solid line; females = dotted line.  
 —○—E —○—VO<sub>2</sub>

**Figure 1.** Mean minute energy expenditure (kcal min<sup>-1</sup>) and oxygen consumption rates (l min<sup>-1</sup>) for aggregate daytime activities of normal-weight males and females as a function of age.


individual values for VAa and  $VD_{physa}/VTa$  ratios, which better correspond to normal distributions (data not shown in tables). Values for AVODa associated with VO<sub>2a</sub> values were found to vary from  $0.059 \pm 0.003$  to  $0.073 \pm 0.004$  ml of O<sub>2</sub> ml<sup>-1</sup> of blood (Table 1). Values for  $VD_{physa}/VTa$  ratios that were calculated based on simultaneous  $VD_{physa}$  and  $VTa$  measurements for healthy subjects free from cardiac and pulmonary diseases (Table 2) correspond to VAa/VEa ratios (i.e.  $1-VD_{physa}/VTa$ ) varying from  $0.619 \pm 0.018$  to  $0.706 \pm 0.032$ .

A 25% reduction in sleep duration for 60% of overweight/obese children, 35% of overweight adults and 55% of their obese counterparts decreased VO<sub>2a</sub>, Qa, VEa and VAa values of the entire cohorts by only 0.5% in boys, 0.6–0.7% in girls and 1.2 and 1.0% in adult males and females, respectively, while VA/Qa ratios were not altered (data not presented in tables).


Maximum mean errors associated with H<sub>p</sub> (–1%), BEE (+2%), ECG and TDEE values (+3.3%) resulted, when combined, in increasing Ea, VO<sub>2a</sub>, Qa, VEa and VAa values by –2.8 to +3.9%. An inverse scenario was observed with minimum mean errors for H<sub>p</sub> (–2%), BEE (+1), ECG and TDEE (–1.0%) values, affecting Ea, VO<sub>2a</sub>, Qa, VEa and VAa values by –1.9 to +4.0%. Variations of H<sub>p</sub>, BEE, TDEE and ECG values did not alter the magnitude of the VAa/Qa ratios (data not given in tables).

## DISCUSSION

The respiratory and cardiovascular parameters determined in the present study are consistent with the range of published values. VAa/Qa ratios reported in this study (or VAa and Qa values) are in agreement with the known values in subjects in the upright




**Figure 2.** Mean minute energy expenditure ( $\text{kcal kg}^{-1} \text{min}^{-1}$ ) and oxygen consumption rates ( $\text{l kg}^{-1} \text{min}^{-1}$ ) for aggregate daytime activities of normal-weight males and females as a function of age.




**Figure 3.** Mean minute energy expenditure ( $\text{kcal m}^{-2} \text{min}^{-1}$ ) and oxygen consumption rates ( $\text{l m}^{-2} \text{min}^{-1}$ ) for aggregate daytime activities of normal-weight males and females as a function of age.

position when their experimental  $VO_2$  demands are within the span of  $VO_2\alpha$  values. This concordance is reflective of the adequacy of the processes and sets of input parameters used for the determination of  $VA\alpha$  and  $Q\alpha$  values (i.e.  $VE\alpha$ ,  $VD_{\text{physa}}/VTA$  and  $VO_2\alpha$ ,  $AVOD\alpha$  respectively), and of course for the calculation of  $VE\alpha$  (i.e.  $E\alpha$ ,  $H_p$ ,  $VQ\alpha$ ) as well as  $VO_2\alpha$  (i.e.  $E\alpha$ ,  $H_p$ ). For instance, mean and individual  $VA/Q$  ratios reported in the literature for resting adults range from  $0.74 \pm 0.09$  to  $0.87 \pm 0.28$  ( $n = 77$ ) and from 0.58 to 1.13 ( $n = 20$ ), respectively (Farhi and Rahn, 1955; West and Dollery, 1960; West, 1962; Lenfant, 1963; Ayres *et al.*,

1964; Johnson and Miller, 1968; West *et al.*, 1974; Zwart *et al.*, 1976; Frostell *et al.*, 1983; Rhodes *et al.*, 1989; Yem *et al.*, 2006). The span of these ratios is in accordance with values of the gap between the 2.5th and 10th percentile  $VA\alpha/Q\alpha$  ratios varying from 0.78 to 1.09 for individuals aged 16.5–<96 years with associated  $VO_2\alpha$  values (0.157–0.426  $\text{l min}^{-1}$ ); this matches well with typical published  $VO_2$  demands (0.236–0.454  $\text{l min}^{-1}$ ) for resting subjects ( $n = 46$ ) aged 19–81 years (Damato *et al.*, 1966; Bachofen *et al.*, 1973). Spans of  $VA\alpha/Q\alpha$  ratios from the 2.5th to 99th percentile in individuals aged 16.5–<25 years and from 10th




**Figure 4.** Mean minute ventilation rates, alveolar ventilation rates and cardiac outputs ( $\text{l min}^{-1}$ ) for aggregate daytime activities of normal-weight males and females as a function of age.



**Figure 5.** Mean minute ventilation rates, alveolar ventilation rates and cardiac outputs ( $\text{l kg}^{-1} \text{min}^{-1}$ ) for aggregate daytime activities of normal-weight males and females as a function of age.

to 99th percentile in those 35–<45 years of age range from 0.91 to 2.14 ( $VEa$  from 9.22 to 30.41) and from 0.92 to 2.40 ( $VEa$  from 9.28 to 31.39  $\text{l min}^{-1}$ ) respectively. By comparison,  $VA/Q$  ratios vary from 0.90 to 2.45 based on  $VA$  and  $Q$  values measured in females aged 20–30 years ( $n=8$ ) inhaling about the same volume of air varying from 9 to 31  $\text{l min}^{-1}$  (Olfert *et al.*, 2004). These 99th percentile  $VAa/Qa$  ratios of 2.14 and 2.40 for  $VEa$  of 30.41 and 31.39  $\text{l min}^{-1}$ , respectively, are also consistent with the higher value of  $VA/Q$  ratio of 2.61 resulting from measurements in men aged 20–30 years ( $n=7$ ) when they were performing activities requiring the higher  $VE$  value of 38.2  $\text{l min}^{-1}$  (Olfert *et al.*, 2004). The 99th percentile  $VAa/Qa$  ratio of 1.93 for  $VO_2a$  ranging from

0.648–0.796  $\text{l min}^{-1}$  in individuals aged 25–<35 year is confirmed by  $VA/Q$  ratios ranging from 2.00 to 2.01 based on simultaneous  $VA$  and  $Q$  measurements in females aged 23.6–30.2 years ( $n=17$ ) by Hopkins *et al.* (2000) during slightly higher  $VO_2$  demands (0.79–0.83  $\text{l min}^{-1}$ ). No published  $VAa/Qa$  ratio is available for older individuals. However,  $VEa$  values used to calculate  $VAa$  values as well as  $Qa$  values for older individuals are in agreement with published values. Spans of  $VEa$  values in males and females aged 45–<65 years between the 2.5th and 99th percentile range from 6.78 to 28.06  $\text{l min}^{-1}$  ( $VO_2a$  from 0.240 to 0.671  $\text{l min}^{-1}$ ), while those in males 65–96 years old vary from 4.52 to 25.16  $\text{l min}^{-1}$  ( $VO_2a$  from 0.157 to 0.611  $\text{l min}^{-1}$ ). Such  $VEa$  values are in



**Figure 6.** Mean minute ventilation rates, alveolar ventilation rates and cardiac outputs ( $\text{l m}^{-2} \text{ min}^{-1}$ ) for aggregate daytime activities of normal-weight males and females as a function of age.

accordance with published  $VE$  values varying from 5.6 to 32.3  $\text{l min}^{-1}$  ( $VO_2$  from 0.236 to 0.797  $\text{l min}^{-1}$ ) in adults aged 45–63 years ( $n=40$ ) and from 5.71 to 25.1  $\text{l min}^{-1}$  ( $VO_2$  from 0.167 to 0.673  $\text{l min}^{-1}$ ) in males 65–91 years old ( $n=29$ ) respectively (Robinson, 1938; Cohn *et al.*, 1954; Tenney and Miller, 1956; Raine and Bishop, 1963; Damato *et al.*, 1966; Bachofen *et al.*, 1973; Nery *et al.*, 1982; Frostell *et al.*, 1983). The span between the 25th and 99th percentile  $Qa$  values for individuals aged 45–96 years ranging from 3.89 to 11.65  $\text{l min}^{-1}$  ( $VO_2a$  from 0.230 to 0.671  $\text{l min}^{-1}$ ) agrees with published values ranging from 3.7 to 12.30  $\text{l min}^{-1}$  ( $VO_2$  from 0.202 to 0.647  $\text{l min}^{-1}$ ) for subjects aged 45–73 years (Reeves *et al.*, 1961; Damato *et al.*, 1966; Emirgil *et al.*, 1967; McGuire *et al.*, 2001;  $n=48$ ).

Regarding children aged 10–<16.5 years, the 2.5th percentile  $VAa/Qa$  ratio of 1.12 ( $VO_2a$  from 0.229 to 0.266  $\text{l min}^{-1}$ ) is in agreement with those varying from 1.07 to 1.17 estimated on the basis of ratios varying from 0.85 to 0.93 for boys aged 11 to 13 years ( $n=9$ ) in the supine position during  $VO_2$  demands ranging from 0.24 to 0.25  $\text{l min}^{-1}$  (Koch and Eriksson, 1973). The latter ratios were increased by 25.3% in order to compensate for the proportional decrease of blood flow that is observed when subjects change from a supine to an upright position (Reeves *et al.*, 1961; Damato *et al.*, 1966; Hossack and Bruce, 1982; Gisolf *et al.*, 2003). Our 99th percentile  $VAa/Qa$  ratio value of 2.16 for the same age groups ( $VO_2a$  from 0.681 to 0.806  $\text{l min}^{-1}$ ) is consistent with higher  $VA/Q$  ratio of 2.49 measured for boys ( $n=9$ ) in the sitting position during much higher  $VO_2$  requirements of 1.14  $\text{l min}^{-1}$  (Koch and Eriksson, 1973). The gap between these lower and upper limits of  $VA/Q$  ratios varying from 1.07 to 2.49 based on data reported in Koch and Eriksson (1973) confirms the magnitude of the span between the 2.5th and 99th percentile  $VAa/Qa$  ratios ranging from 1.12 to 2.16 in children aged 10–<16.5 years. The magnitudes of  $VAa$  and  $Qa$  values for younger children are confirmed by published measurements. For instance, the span between the 25th and 90th percentile  $VAa$  values ranges from 5.26 to 8.63 ( $VO_2a$  from 0.259 to 0.417  $\text{l min}^{-1}$ ) in children aged 7–<10 years. By comparison,  $VA$  values varying from 5.03 to 9.03  $\text{l min}^{-1}$  have

been measured in those aged 6–17 years ( $n=56$ ) during comparable  $VO_2$  demands ranging from 0.262 to 0.389  $\text{l min}^{-1}$  (Zapletal *et al.*, 1987). The 97.5th percentile  $Qa$  values of 6.73  $\text{l min}^{-1}$  in boys and 6.23  $\text{l min}^{-1}$  in girls aged 7–<10 years for  $VO_2a$  of 0.487 and 0.446  $\text{l min}^{-1}$ , respectively, are also in accordance with mean values of 6.8  $\text{l min}^{-1}$  in males ( $n=12$ ) and 6.60  $\text{l min}^{-1}$  in females ( $n=12$ ) aged 7–9 years measured during light activities with  $VO_2$  demands of 0.55 and 0.51  $\text{l min}^{-1}$  respectively (Turley and Wilmore, 1997a).

As expected, mean  $VAa/Qa$  ratios in children aged 5–<16.5 years ( $1.49 \pm 0.12$  to  $1.53 \pm 0.24$ ) are higher than those for older individuals 16.5–96 years old ( $1.22 \pm 0.27$  to  $1.36 \pm 0.28$ ). In response to higher oxygen demands associated with higher energy expenditures in children aged 5–<16.5 years ( $0.044 \pm 0.014$  to  $0.063 \pm 0.014 \text{ kcal kg}^{-1} \text{ min}^{-1}$ ,  $1.40 \pm 0.40$  to  $1.72 \pm 0.56 \text{ kcal m}^{-2} \text{ min}^{-1}$ ), when compared with lower oxygen demands in older individuals aged 16.5–96 years ( $0.025 \pm 0.007$  to  $0.038 \pm 0.007 \text{ kcal kg}^{-1} \text{ min}^{-1}$ ,  $0.88 \pm 0.25$ – $1.44 \pm 0.25 \text{ kcal m}^{-2} \text{ min}^{-1}$ ),  $VA$  (and thus  $VE$ ) values increase in order to sustain adequate oxygen blood concentrations, while the  $Q$  values rise in order to increase oxygen transport to all body tissues. Higher oxygen uptakes in children compared with those in older individuals are reflected by higher number of alveoli per unit of body weight and body surface area. For instance, the number of alveoli determined in children 4 and 8 years of age is  $15.86$  and  $11.20 \times 10^6 \text{ alveoli kg}^{-1}$ , or  $383.6$  and  $304.4 \times 10^6 \text{ alveoli m}^{-2}$  respectively, compared with much lower values in adults:  $3.84 \times 10^6 \text{ alveoli kg}^{-1}$  or  $155.8 \times 10^6 \text{ alveoli m}^{-2}$  (Dunnill, 1962). These values are consistent with those reported by Davies and Reid (1970) as well as Angus and Thurlbeck (1972).

Thus, alveolar ventilation rates must maintain a relatively high level of alveolar and arterial oxygen partial pressure in order to compensate for temporary biochemical differences that are observed in children aged 5–<16.5 years compared with older individuals. Lower blood hemoglobin concentrations and slightly higher concentrations of 2,3-diphosphoglycerate are observed in children 5–<10 years old compared with those aged 10–<16.5 years (Motoyama *et al.*, 1990). Higher concentrations of 2,3-diphosphoglycerate in red cells increase the oxygen



unloading from hemoglobin at the tissue level (Osaki and Delivoria-Papadopoulos, 1970; Card and Brain, 1973; Osaki, 1973; Motoyama *et al.*, 1974). Mean hemoglobin levels for children aged 2–5, 6–8, 10–12 and 14–16 years are  $11.9 \pm 1.2$  ( $n=22$ ),  $12.6 \pm 0.8$  ( $n=41$ ),  $13.2 \pm 0.9$  ( $n=54$ ) and  $14.4 \pm 1.4$  g dl<sup>-1</sup> ( $n=34$ ) in boys and  $12.4 \pm 0.9$  ( $n=20$ ),  $12.7 \pm 1.0$  ( $n=10$ ),  $13.2 \pm 1.0$  ( $n=29$ ) and  $13.4 \pm 1.2$  g dl<sup>-1</sup> ( $n=15$ ) in girls, respectively (Spurr *et al.*, 1992). These values, which are in agreement with other blood hemoglobin concentrations varying from  $12.99 \pm 0.31$  to  $13.9 \pm 1.3$  g dl<sup>-1</sup> ( $n=186$ ) for children 7–13.7 years of age (Åstrand, 1952; Eriksson *et al.*, 1971; Koch and Eriksson, 1973; Turley and Wilmore, 1997a, b; Obert *et al.*, 2003; Vinet *et al.*, 2003), are lower than those for adults aged 18–89 years ( $n=504$ ) ranging from  $13.00 \pm 1.25$  to  $15.9 \pm 1.2$  g dl<sup>-1</sup> (Rotta *et al.*, 1956; Tenney and Miller, 1956; Åstrand *et al.*, 1964; Ekblom *et al.*, 1968; Holmér *et al.*, 1974; Kanstrup and Ekblom, 1982; Bebout *et al.*, 1989; Stringer *et al.*, 1997; Proctor *et al.*, 1998a, b, 2003; Sun *et al.*, 2000; Poole *et al.*, 2002; Mourtzakis *et al.*, 2004; Beck *et al.*, 2006). Overall, immature mechanisms for oxygen transport to body tissues with higher energy expenditures in children 5–<16.5 years old provide a reasonable explanation for the unique values of VA and Q specific to this age group.

The magnitude of inter-individual variability of 8.4 for cardiac output and 13.4 for alveolar ventilation rate was calculated as the ratio of the highest 99th percentiles of 0.330 and 0.553 l kg<sup>-1</sup> per day (Tables 6 and 7) to the lowest 1st percentiles of 0.039 and 0.041 l kg<sup>-1</sup> per day, respectively (data not shown in tables) in males and females aged 5–96 years. The magnitude of human variability in Q and VA values, as reflected by the lowest 50th percentiles of 0.084 and 0.100 l kg<sup>-1</sup> per day (Tables 6 and 7) and the highest 95th percentiles of 0.262 and 0.413 l kg<sup>-1</sup> per day (data not shown in tables) correspond to factors of 3.1 and 4.1, respectively. The impact of such inter-individual variability in Q (i.e. 3.1–8.4) and VA values (i.e. 4.1–13.4) should be assessed along with the variability in other pharmacokinetic determinants, in order to evaluate the adequacy of the default uncertainty factor or the human kinetic adjustment factor currently used in health risk assessment (Renwick, 2000; World Health Organization, 2005).

## CONCLUSION

The present study provides a complete and original set of key respiratory and cardiovascular parameters (i.e.  $E_a$ ,  $AVOD_a$ ,  $VO_{2a}$ ,  $VE_a$ ,  $Q_a$ ,  $VA_a$ , values and  $VD_{physa}/VT_a$ ,  $VA_a/Q_a$  ratios), with their distributions, for healthy normal-weight males and females aged 5–96 years old during their aggregate daytime activities. As done by Brochu *et al.* (2011) for the selection of input literature data when calculating  $H_p$  and  $VQ_a$  values, solely data measured in subjects in the upright position during  $VO_2$  demands that were within the span of  $VO_{2a}^*$  values were used in this study. Such a procedure assures that data included in the calculation processes of  $VO_{2a}$ ,  $VE_a$ ,  $Q_a$ ,  $VA_a$ , values and  $VA_a/Q_a$  ratios adequately describe daytime activities for individuals of different age groups. The fact that the spans of  $VO_{2a}$  values per age group appear to be in agreement with those for  $VO_{2a}^*$  provides added value to this approach.

Determination of energy expenditures during aggregate daytime activities (i.e.  $E_a$ ) for each age group by subtracting published BEE from TDEE values that are measured for the same subjects by the DLW method is unique. Indirect calorimetry

measurements ( $n=902$ ) in normal-weight males and females and disappearance rates of oral doses of water isotopes ( $^2H_2O$  and  $H_2^{18}O$ ) in urine for an aggregate period of over 14 000 days were used for the calculation of  $E_a$  values. In addition, the accuracy of  $VO_{2a}$ ,  $VE_a$ ,  $Q_a$ ,  $VA_a$ , values expressed in l min<sup>-1</sup>, l kg<sup>-1</sup> min<sup>-1</sup> as well as l m<sup>-2</sup> min<sup>-1</sup> and  $VA_a/Q_a$  ratios is enhanced by the facts that: (1) the weight and height, as well as the BEE and TDEE values used in the calculation processes, were available for each subject when conducting the DLW method; (2) each TDEE value systematically encompasses voluntary and involuntary energy expended in unrestrained free-living subjects each minute of the day, 24 h per day, on a daily basis during 7–21 days; and (3) in the worst case scenario, simultaneous extreme mean errors for  $H_p$  (–2 to –1%), BEE (+1 to +2%) and TDEE (–1.0 to +3.3%) values only affect  $E_a$ ,  $VO_{2a}$ ,  $Q_a$ ,  $VE_a$ ,  $VA_a$  values by –2.8 to +4.0%.

The absorption rates of inhaled gases and vapors of xenobiotics with high or low blood/gas phase solubility ratios are increased by higher VA or Q values respectively (Klaassen, 1996). In the present study, generally higher 2.5nd to 99th percentile  $VE_a$  (0.132–0.774 l kg<sup>-1</sup> min<sup>-1</sup>, 4.42–21.69 l m<sup>-2</sup> min<sup>-1</sup>),  $VA_a$  (0.093–0.553 l kg<sup>-1</sup> min<sup>-1</sup>, 3.09–15.53 l m<sup>-2</sup> min<sup>-1</sup>) and  $Q_a$  values (0.065–0.330 l kg<sup>-1</sup> min<sup>-1</sup>, 2.17–9.46 l m<sup>-2</sup> min<sup>-1</sup>), as well as  $VA_a/Q_a$  ratios (1.12–2.16) were found in normal-weight children 5–<16.5 years of age when compared with older individuals ( $VE_a$  0.076–0.461 l kg<sup>-1</sup> min<sup>-1</sup>, 2.80–16.99 l m<sup>-2</sup> min<sup>-1</sup>;  $Q_a$  0.045–0.201 l kg<sup>-1</sup> min<sup>-1</sup> and 1.63–7.24 l m<sup>-2</sup> min<sup>-1</sup>;  $VA_a$  0.047–0.312 l kg<sup>-1</sup> min<sup>-1</sup> and 1.73–11.63 l m<sup>-2</sup> min<sup>-1</sup>;  $VA_a/Q_a$  ratios 0.78–2.40). Therefore, all factors being equal, the age-related differences in the respiratory rates and cardiac output can have a direct effect on the intake and uptake of inhaled gases and vapors, notably liposoluble air pollutants by the respiratory tract in younger individuals.

## Acknowledgments

Appreciation is expressed to Mr André Thibault for his participation in the review of published data regarding the variation of sleep duration as a function of body mass index in individuals. The authors wish to thank Mr John Francis Niederreiter for the linguistic revision of this manuscript. The authors wish to point out that the views expressed in this paper may not reflect the official policy of the Québec Ministry of Sustainable Development, Environment and Parks.

## Declaration of Interest

The authors declare that there are no conflicts of interest.

## REFERENCES

- Allen CJ, Jones NL, Killian KJ. 1984. Alveolar gas exchange during exercise: a single-breath analysis. *J. Appl. Physiol.* **57**: 1704–1709.
- Angus GE, Thurlbeck WM. 1972. Number of alveoli in the human lung. *J. Appl. Physiol.* **32**(4): 483–485.
- Arms AD, Travis CC. 1988. *Physiological Parameters in Pharmacokinetic Modeling*. US Environmental Protection Agency 600/6-88/004. Final report. (NTIS PB 88-196019). Washington, DC.
- Åstrand P-O. 1952. *Experimental Studies of Physical Working Capacity in Relation to Sex and Age*. Kungliga Gymnastiska Central Institutet: Stockholm/Ejnar Munksengaard: Copenhagen.
- Åstrand P-O, Cuddy TE, Saltin B, Stenberg J. 1964. Cardiac output during submaximal and maximal work. *J. Appl. Physiol.* **19**: 268–274.
- Ayres SM, Criscitiello A, Grabovsky E. 1964. Components of alveolar-arterial O<sub>2</sub> difference in normal man. *J. Appl. Physiol.* **19**: 43–47.

- Bachofen H, Hobi HJ, Scherrer M. 1973. Alveolar-arterial  $N_2$  gradients at rest and during exercise in healthy men of different ages. *J. Appl. Physiol.* **34**: 137-142.
- Baudry PH, Wise MB, Seely JE. 1966. Respiratory gas exchange at rest and during exercise in normal and asthmatic children. *Am. Rev. Respir. Dis.* **95**: 248-254.
- Bebout DE, Story D, Roca J, Hogan MC, Poole DC, Camarena RG, Ueno O, Haab P, Wagner PD. 1989. Effects of altitude acclimatization on pulmonary gas exchange during exercise. *J. Appl. Physiol.* **67**(6): 2286-2295.
- Beck KC, Randolph LN, Bailey KR, Wood CM, Snyder EM, Johnson BD. 2006. Relationship between cardiac output and oxygen consumption during upright cycle exercise in healthy humans. *J. Appl. Physiol.* **101**(5): 1474-1480.
- Becklake MR, Varvis CJ, Pengelly LD, Kenning S, McGregor M, Bates DV. 1962. Measurement of pulmonary blood flow during exercise using nitrous oxide. *J. Appl. Physiol.* **17**: 579-586.
- Bernsteins MS, Costanza MC, Morabia A. 2001. Physical activity of urban adults: a general population survey in Geneva. *Soz. Praventivmed.* **46**: 049-059.
- Bohr C. 1891. Ueber die Lungenathmung. *Skand. Arch. Physiol.* **2**: 236-268.
- Brochu P, Ducré-Robitaille J-F, Brodeur J. 2006a. Physiological daily inhalation rates for free-living individuals aged 1 month to 96 years, using data from doubly labeled water measurements: a proposal for air quality criteria, standard calculations and health risk assessment. *Hum. Ecol. Risk Assess.* **12**(4): 675-701.
- Brochu P, Ducré-Robitaille J-F, Brodeur J. 2006b. Physiological daily inhalation rates for free-living pregnant and lactating adolescents and women aged 11 to 55 years, using data from doubly labeled water measurements for use in health risk assessment. *Hum. Ecol. Risk Assess.* **12**(4): 702-735.
- Brochu P, Ducré-Robitaille J-F, Brodeur J. 2006c. Physiological daily inhalation rates for free-living individuals aged 2.6 months to 96 years based on doubly labeled water measurements: comparison with time-activity-ventilation and metabolic energy conversion estimates. *Hum. Ecol. Risk Assess.* **12**(4): 736-761.
- Brochu P, Brodeur J, Krishnan K. 2011. Derivation of physiological inhalation rates in children, adults and elderly based on nighttime and daytime respiratory parameters. *Inhal. Toxicol.* **23**(2): 74-94.
- Bursztin S, Elwyn D, Askanazi J, Kinney J. 1989. *Energy Metabolism, Indirect Calorimetry, and Nutrition*. Williams and Wilkins: Baltimore, MD.
- Card RT, Brain MC. 1973. The anemia of childhood: evidence for a physiologic response to hyperphosphatemia. *New Engl. J. Med.* **288**: 388.
- Cohn JE, Carroll DG, Armstrong BW, Shepard RH, Riley RL. 1954. Maximal diffusing capacity of the lung in normal male subjects of different ages. *J. Appl. Physiol.* **6**: 588-597.
- Cook CD, Cherry D, O'Brien RB, Karlberg P, Smith CA. 1955. Studies of respiratory physiology in the newborn infant. i. observation on normal premature and full-term infant. *J. Clin. Invest.* **34**: 975-982.
- Craig DB, Wahba MW, Don HF, Couture JG, Becklake MR. 1971. 'Closing volume' and its relationship to gas exchange in seated and supine positions. *J. Appl. Physiol.* **31**: 717-721.
- Dagenais GR, Oriol A, McGregor M. 1966. Hemodynamic effects of carbohydrate and protein meals in man: rest and exercise. *J. Appl. Physiol.* **21**: 1157-1162.
- Damato AN, Galante JG, Smith WM. 1966. Hemodynamic response to treadmill exercise in normal subjects. *J. Appl. Physiol.* **21**: 959-966.
- Davies G, Reid L. 1970. Growth of the alveoli and pulmonary arteries in childhood. *Thorax* **25**: 669-681.
- Dempsey JA, Hanson PG, Henderson KS. 1984. Exercise-induced arterial hypoxaemia in healthy human subjects at sea level. *J. Physiol.* **355**: 161-175.
- Dibski DW, Smith DJ, Jensen R, Norris SR, Ford GT. 2005. Comparison and reliability of two non-invasive acetylene uptake techniques for the measurement of cardiac output. *Eur. J. Appl. Physiol.* **94**: 670-680.
- Donevan RE, Anderson NM, Sekelj P, Papp O, McGregor M. 1962. Influence of voluntary hyperventilation on cardiac output. *J. Appl. Physiol.* **17**: 487-491.
- Dunnill MS. 1962. Postnatal growth of the lung. *Thorax* **17**: 329-333.
- Durnin JVGA, Passmore R. 1967. *Energy, Work and Leisure*. Heinemann Educational Books, London; 24-105.
- Eisenmann JC, Ekkekakis P, Holmes M. 2006. Sleep duration and overweight among Australian children and adolescents. *Acta Paediat.* **95**(9): 956-963.
- Eklom B, Astrand PO, Saltin B, Stenberg J, Wallstrom B. 1968. Effect of training on circulatory response to exercise. *J. Appl. Physiol.* **24**: 518-528.
- Elia M. 1992. Organ and tissue contribution to metabolic rate. In *Energy Metabolism: Tissue Determinants and Cellular Corollaries*, Kenney JM, Tucker HN (eds). Lippincott-Raven: New York; 61-79.
- Elia M. 1997. Tissue distribution and energetics in weight loss and undernutrition. In *Physiology. Stress and Malnutrition: Functional Correlates, Nutritional Intervention*, Kenney JM, Tucker HN (eds). Lippincott-Raven: New York; 383-411.
- Emirgil C, Sobol BJ, Campodonico S, Herbert WH, Mechkati R. 1967. Pulmonary circulation in the aged. *J. Appl. Physiol.* **23**: 631-640.
- Enghoff H. 1938. Volumen inefficax, Bemerkungen zur Frage des schädlichen Raumes. *Upsala Lakareforen Forth* **44**: 191-218.
- Eriksson BO, Grimby G, Saltin B. 1971. Cardiac output and arterial blood gases during exercise in pubertal boys. *J. Appl. Physiol.* **31**: 348-352.
- Farhi LE, Rahn H. 1955. Gas stores of the body and the unsteady state. *J. Appl. Physiol.* **7**: 472-484.
- Ferrannini E. 1988. The theoretical basis of indirect calorimetry: a review. *Metabolism* **37**: 287-301.
- Fick A. 1870. Über die messung des blutquantums in den hertzventrikel. (On the measurement of blood mass in the heart ventricles). *Sitz ber Physik-Med Ges Wurzburg* **2**: 16-28.
- Fowler WS. 1948. Lung function studies. II: The respiratory dead space. *Am. J. Physiol.* **154**: 406-416.
- Folkow B, Pappenheimer JR. 1955. Components of the respiratory dead space and their variation with pressure breathing and with bronchoactive drugs. *J. Appl. Physiol.* **8**: 102-110.
- Frick MH, Somer T. 1964. Base-line effects on response of stroke volume to leg exercise in the supine position. *J. Appl. Physiol.* **19**: 639-643.
- Froeb HF. 1962. Stimulation of ventilation in emphysema by passively induced body motion. *J. Appl. Physiol.* **17**: 771-774.
- Frostell C, Pande JN, Hedenstierna G. 1983. Effects of high-frequency breathing on pulmonary ventilation and gas exchange. *J. Appl. Physiol.* **55**: 1854-1861.
- Gisolf J, Wilders R, Immink RV, van Lieshout JJ, Karemaker JM. 2003. Tidal volume, cardiac output and functional residual, capacity determine end-tidal  $CO_2$  transient during standing up in humans. *J. Physiol.* **554**(2): 579-590.
- Guyton AC. 1991. *Textbook of Medical Physiology*, 8th edn. W.B. Saunders/Harcourt Brace Jovanovich: Philadelphia, PA.
- Haddad S, Charest-Tardif G, Tardif R. 2006. Developpement of physiologically based toxicokinetic models for improving the human indoor exposure assessment to water contaminants: trichloroethylene and trihalomethanes. *J. Toxicol. Environ. Health Part A* **69**: 2095-2136.
- Hermansen L, Eklom B, Saltin B. 1970. Cardiac output during submaximal and maximal treadmill and bicycle exercise. *J. Appl. Physiol.* **29**: 82-86.
- Holmér I, Stein EM, Saltin B, Eklom B, Astrand PO. 1974. Hemodynamic and respiratory responses compared in swimming and running. *J. Appl. Physiol.* **37**: 49-54.
- Hopkins SR, Barker RC, Brutsaert TD, Gavin TP, Olfert IM, Veisel S, Wagner PD. 2000. Pulmonary gas exchange during exercise in women: effects of exercise type and work increment. *J. Appl. Physiol.* **89**: 721-730.
- Hossack KF, Bruce RA. 1982. Maximal cardiac function in sedentary normal men and women: comparison of age-related changes. *J. Appl. Physiol.* **53**: 799-804.
- International Dietary Energy Consultancy Group. 1990. *The Doubly-labelled Water Method for Measuring Energy Expenditure: a Consensus Report by the IDECG Working Group. Technical Recommendation for Use in Humans*, Prentice AM (ed.). NAHRES-4. IAEA: Vienna. Available at: <http://www.unu.edu/unupress/food2/UID05E/UID05E00.HTM>
- Institute of Medicine. 2002. Appendix I: doubly labeled water data used to predict energy expenditure. In *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids* (Macronutrients). Food and Nutrition Board. National Academies Press: Washington, DC; 1104-1202. Available at: <http://books.nap.edu/books/0309085373/html/index.html>

- Jones WB, Finchum RN, Russell RO Jr, Reeves TJ. 1970. Transient cardiac output response to multiple levels of supine exercise. *J. Appl. Physiol.* **28**: 183–189.
- Johnson RL Jr, Miller JM. 1968. Distribution of ventilation, blood flow, and gas transfer coefficients in the lung. *J. Appl. Physiol.* **25**: 1–15.
- Johnson RL Jr, Spicer WS, Bishop JM, Forster RE. 1960. Pulmonary capillary blood volume, flow and diffusing capacity during exercise. *J. Appl. Physiol.* **15**: 893–902.
- Kanstrup IL, Ekblom B. 1978. Influence of age and physical activity on central hemodynamics and lung function in active adults. *J. Appl. Physiol.* **45**: 709–717.
- Kanstrup IL, Ekblom B. 1982. Acute hypervolemia, cardiac performance, and aerobic power during exercise. *J. Appl. Physiol.* **52**: 1186–1191.
- Kerr AA. 1976. Dead space ventilation in normal children and children with obstructive airways disease. *Thorax* **31**: 63–69.
- Klaassen CD. 1996. *Casarett and Doull's Toxicology: The Basic Science of Poisons*, 5th edn, Klaassen CD, Amdur MO, Doull J (eds). McGraw-Hill: New York.
- Koch G, Eriksson BO. 1973. Effect of physical training on pulmonary ventilation and gas exchange during submaximal and maximal work in boys aged 11 to 13 years. *Scand. J. Clin. Lab. Invest.* **31**: 87–94.
- Krishnan K, Andersen ME. 2001. Physiologically-based pharmacokinetic modeling in toxicology. In *Principles and Methods in Toxicology*, Hayes W (ed.). Taylor & Francis: New York; 193–141.
- Krone RJ, Goldbarg AN, Balkoura M, Schuessler R, Resnekov L. 1972. Effects of cigarette smoking at rest and during exercise. II. Role of venous return. *J. Appl. Physiol.* **32**: 745–748.
- Lenfant C. 1963. Measurement of ventilation/perfusion distribution with alveolar-arterial differences. *J. Appl. Physiol.* **18**: 1090–1094.
- Lewis SF, Taylor WF, Graham RM, Pettinger WA, Schutte JE, Blomqvist CG. 1983. Cardiovascular responses to exercise as functions of absolute and relative work load. *J. Appl. Physiol.* **54**: 1314–1323.
- Malmberg R. 1966. Pulmonary gas exchange at exercise and different body postures in man. *Scand. J. Resp. Dis.* **47**: 92–102.
- McGuire DK, Levine BD, Williamson JW, Snell PG, Blomqvist CG, Saltin B, Mitchell JH. 2001. A 30-year follow-up of the Dallas bed rest and training study: I. Effect of age on the cardiovascular response to exercise. *Circulation* **104**(12): 1350–1357.
- Mellergaard K. 1966. The alveolar-arterial oxygen difference: its size and components in normal man. *Acta Physiol. Scand.* **67**: 10–20.
- Miyamoto Y, Niizeki K, Kawahara K, Doi K. 1989. Cardiodynamic factors affecting hyperpnea during steady-state exercise in man. *Jpn. J. Physiol.* **39**(3): 411–420.
- Mosteller RD. 1987. Simplified calculation of body surface area. *New Engl. J. Med.* **317**(17): 1098.
- Motoyama EK, Zigas CJ, Troll G. 1974. Functional basis of childhood anemia (abstract). *Am. Soc. Anesth.* **41**: 283.
- Motoyama EK, Davis PJ, Lewis Cohn E. 1990. Chapter 2: Respiratory physiology in infants and children. In *Smith's Anesthesia for Infants and Children*, 5th edn. Mosby: St Louis; 11–76.
- Mourtzakis M, Gonzalez-Alonso J, Graham TE, Saltin B. 2004. Hemodynamics and O<sub>2</sub> uptake during maximal knee extensor exercise in untrained and trained human quadriceps muscle: effects of hyperoxia. *J. Appl. Physiol.* **97**(5): 1796–1802.
- Nelson NM, Prod'homme LS, Cherry RB, Lipsitz PJ, Smith CA. 1962. Pulmonary functions in the newborn infant. I. Methods: ventilation and gaseous metabolism. *Pediatrics* **31**: 963–973.
- Nery LE, Wasserman K, Andrews JD, Huntsman DJ, Hansen JE, Whipp BJ. 1982. Ventilatory and gas exchange kinetics during exercise in chronic airways obstruction. *J. Appl. Physiol.* **53**: 1594–1602.
- Nottin S, Vinet A, Stecken F, Nguyen LD, Ounissi F, Lecoq AM, Obert P. 2002. Central and peripheral cardiovascular adaptations during a maximal cycle exercise in boys and men. *Med. Sci. Sports Exerc.* **34**(3): 456–463.
- Obert P, Mandigouts S, Nottin S, Vinet A, N'Guyen LD, Lecoq AM. 2003. Cardiovascular responses to endurance training in children: effect of gender. *Eur. J. Clin. Invest.* **33**(3): 199–208.
- Olfert IM, Balouch J, Kleinsasser A, Knapp A, Wagner H, Wagner PD, Hopkins SR. 2004. Does gender affect human pulmonary gas exchange during exercise? *J. Physiol.* **557**(2): 529–541.
- Oski FA. 1973. Designation of anemia on a functional basis. *J. Pediatr.* **83**: 353.
- Oski FA, Delivoria-Papadopoulos M. 1970. The red cell, 2,3-diphosphoglycerate, and tissue oxygen release. *J. Pediatr.* **77**: 941.
- Ouellet Y, Poh SC, Becklake MR. 1969. Circulatory factors limiting maximal aerobic exercise capacity. *J. Appl. Physiol.* **27**: 874–880.
- Pernow B, Saltin B. 1971. Availability of substrates and capacity for prolonged heavy exercise in man. *J. Appl. Physiol.* **31**: 416–422.
- Podolsky A, Eldridge MW, Richardson RS, Knight DR, Johnson EC, Hopkins SR, Johnson DH, Michimata H, Grassi B, Feiner J, Kurdak SS, Bickler PE, Severinghaus JW, Wagner PD. 1996. Exercise-induced VA/Q inequality in subjects with prior high-altitude pulmonary edema. *J. Appl. Physiol.* **81**(2): 922–932.
- Poole JG, Lawrenson L, Kim J, Brown C, Richardson RS. 2002. Vascular and metabolic response to cycle exercise in sedentary humans: effect of age. *Am. J. Physiol. Heart Circul. Physiol.* **284**: H1251–H1259.
- Price K, Haddad S, Krishnan K. 2003. Physiological modeling of age-specific changes of pharmacokinetics of organic chemicals in children. *J. Toxicol. Environ. Health, Pt A* **66**: 417–433.
- Proctor DN, Beck KC, Shen PH, Eickhoff TJ, Halliwill JR, Joyner MJ. 1998a. Influence of age and gender on cardiac output–VO<sub>2</sub> relationships during submaximal cycle ergometry. *J. Appl. Physiol.* **84**(2): 599–605.
- Proctor DN, Shen PH, Dietz NM, Eickhoff TJ, Lawler LA, Ebersold EJ, Loeffler DL, Joyner MJ. 1998b. Reduced leg blood flow during dynamic exercise in older endurance-trained men. *J. Appl. Physiol.* **85**(1): 68–75.
- Proctor DN, Newcomer SC, Koch DW, Le Khoi U, MacLean DA, Leuenberger UA. 2003. Leg blood flow during submaximal cycle ergometry is not reduced in healthy older normally active men. *J. Appl. Physiol.* **94**: 1859–1869.
- Raine JM, Bishop JM. 1963. A difference in O<sub>2</sub> tension and physiological dead space in normal man. *J. Appl. Physiol.* **18**: 284–288.
- Reeves JT, Grover RF, Blount SG Jr, Filley GF. 1961. Cardiac output response to standing and treadmill walking. *J. Appl. Physiol.* **16**: 283–288.
- Renwick AG. 2000. The use of safety or uncertainty factors in the setting of acute reference doses. *Food Addit. Contam.* **17**(7): 627–635.
- Rhodes CG, Valind SO, Brudin LH, Wollmer PE, Jones T, Bickingham PD, Hughes JMB. 1989. Quantification of regional v/Q ratios in humans by use of PET. II. Procedure and normal values. *J. Appl. Physiol.* **66**(4): 1905–1913.
- Rice AJ, Thornton AT, Gore CJ, Scroop GC, Greville HW, Wagner H, Wagner PD, Hopkins SR. 1999. Pulmonary gas exchange during exercise in highly trained cyclists with arterial hypoxemia. *J. Appl. Physiol.* **87**(5): 1802–1812.
- Robinson S. 1938. Experimental studies of physical fitness in relation to age. *Arbeits. Physiol.* **10**: 251–323.
- Rotta A, Cánepa A, Hurtado A, Velásquez T, Chávez R. 1956. Pulmonary circulation at sea level and at high altitudes. *J. Appl. Physiol.* **9**: 328–336.
- Sharma JD, Saxana RK, Rastogi SK. 1977. Cardiac output of Indian men by a non-invasive method. The indirect Fick principle. *Ind. J. Physiol. Pharmac.* **21**(4): 347–352.
- Spurr GB, Reina JC, Hoffmann RG. 1992. Basal metabolic rate of Colombian children 2–16 y of age: ethnicity and nutritional status. *Am. J. Clin. Nutr.* **56**: 623–629.
- Stahlman MT, Meece NJ. 1957. Pulmonary ventilation and diffusion in the human newborn infant. *J. Clin. Invest.* **36**: 1081–1091.
- Stringer WW, Hansen JE, Wasserman K. 1997. Cardiac output estimated noninvasively from oxygen uptake during exercise. *J. Appl. Physiol.* **82**(3): 908–912.
- Sun XG, Hansen JE, Ting H, Chuang ML, Stringer WW, Adame D, Wasserman K. 2000. Comparison of exercise cardiac output by the Fick principle using oxygen and carbon dioxide. *Chest* **118**(3): 631–640.
- Tabakin BS, Hanson JS, Merriam TW Jr, Caldwell EJ. 1964. Hemodynamic response of normal men to graded treadmill exercise. *J. Appl. Physiol.* **19**: 457–464.
- Tenney SM, Miller RM. 1956. Dead space ventilation in old age. *J. Appl. Physiol.* **9**: 321–327.
- Torre-Bueno JR, Wagner PD, Saltzman HA, Gale GE, Moon RE. 1985. Diffusion limitation in normal humans during exercise at sea level and simulated altitude. *J. Appl. Physiol.* **58**(3): 989–995.
- Travis CC, Hattermer-Frey HA. 1991. Physiological pharmacokinetic models. In *Statistics in Toxicology*, Krewski D, Franklin C (eds). Gordon and Breach: New York.
- Turell DJ, Alexander JK. 1964. Experimental evaluation of Weir's formula for estimating metabolic rate in man. *J. Appl. Physiol.* **19**(5): 946–948.
- Turley KR, Wilmore JH. 1997a. Cardiovascular responses to submaximal exercise in 7- to 9-yr-old boys and girls. *Med. Sci. Sports Exerc.* **29**(6): 824–832.



- Turley KR, Wilmore JH. 1997b. Cardiovascular responses to treadmill and cycle ergometer exercise in children and adults. *J. Appl. Physiol.* **83**(3): 948–957.
- US Environmental Protection Agency. 1992. Guideline for exposure assessment; notice. *Fed. Reg.* **57**(104): 22888–22938.
- US Environmental Protection Agency. 1988. Reference physiological parameters in pharmacokinetic modeling. Final report. EPA/600/6-88/004, PB88-19019. USEPA: Washington, DC.
- Valcke M, Krishnan K. 2009. Physiologically based pharmacokinetic modeling in the risk assessment of developmental toxicants. In *Developmental Toxicology*, 3rd edn, Hansen DK, Abbott BD (eds). Informa Healthcare: New York; 243–273.
- Vinet A, Nottin S, Lecoq AM, Obert P. 2002. Cardiovascular responses to progressive cycle exercise in healthy children and adults. *Int. J. Sports Med.* **23**(4): 242–246.
- Vinet A, Mandigout S, Nottin S, Nguyen LD, Lecoq A-M, Courteix D, Obert P. 2003. Influence of body composition, hemoglobin concentration, cardiac size and function of gender differences in maximal oxygen uptake in prepubertal children. *Chest* **124**: 1494–1499.
- Wagner PD, Gale GE, Moon RE, Torre-Bueno JR, Stolp WB, Saltzman HA. 1986. Pulmonary gas exchange in humans exercising at sea level and simulated altitude. *J. Appl. Physiol.* **61**: 260–270.
- Weir JB, de V. 1949. New methods for calculating metabolic rate with special reference to protein metabolism. *J. Physiol.* **109**: 1–9.
- West JB. 1962. Regional differences in gas exchange in the lung of erect man. *J. Appl. Physiol.* **17**(6): 893–898.
- West JB, Dollery CT. 1960. Distribution of blood flow and ventilation-perfusion ratio in the lung, measured with radioactive CO<sub>2</sub>. *J. Appl. Physiol.* **15**: 405–410.
- West JB, Wagner PD, Derks CM. 1974. Gas exchange in distributions of VA–Q ratios: partial pressure–solubility diagram. *J. Appl. Physiol.* **37**: 533–540.
- Whipp BJ, Wasserman K. 1969. Alveolar–arterial gas tension differences during graded exercise. *J. Appl. Physiol.* **27**: 361–365.
- World Health Organization. 2005. *Chemical-specific and Adjustment Factors for Interspecies Differences and Human Variability: Guidance Document for Use of Data in Dose/Concentration–Response Assessment*. Harmonization Project Document No. 2. WHO: Geneva.
- Yem JS, Turner MJ, Baker AB, Young IH, Crawford ABH. 2006. A tidally breathing model of ventilation, perfusion and volume in normal and diseased lungs. *Br. J. Anaesth.* **97**(5): 718–731.
- Zapletal A, Samanek M, Paul T. 1987. *Lung Function in Children and Adolescents. Methods, Reference Values*. H. Herzog Series. Progr. Resp. Res. **22**. Karger: Basel.
- Zeidifard E, Silverman M, Godfrey S. 1972. Reproducibility of indirect (CO<sub>2</sub>) Fick method for calculation of cardiac output. *J. Appl. Physiol.* **33**(1): 141–143.
- Zwart A, Seagrave RC, Van Dieren A. 1976. Ventilation–perfusion ratio obtained by a noninvasive frequency response technique. *J. Appl. Physiol.* **41**: 419–424.